

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number
WO 02/20501 A2

(51) International Patent Classification⁷: **C07D 295/14**

(21) International Application Number: **PCT/EP01/09926**

(22) International Filing Date: 27 August 2001 (27.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00203067.4 4 September 2000 (04.09.2000) EP

(71) Applicant (*for all designated States except US*):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Patent
Department, Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MEERPOEL,**
Lieven [BE/BE]; Janssen Pharmaceutica N.V., Turn-
houtseweg 30, B-2340 Beerse (BE). **ROEVENS, Peter,**
Walter, Maria [BE/BE]; Janssen Pharmaceutica N.V.,
Turnhoutseweg 30, B-2340 Beerse (BE). **BACKX, Leo,**
Jacobus, Jozef [BE/BE]; Janssen Pharmaceutica N.V.,
Turnhoutseweg 30, B-2340 Beerse (BE). **VAN DER**
VEKEN, Louis, Jozef, Elisabeth [BE/BE]; Janssen
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse
(BE). **VIELLEVOYE, Marcel** [NL/BE]; Janssen Phar-
maceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Agent: **VERBERCKMOES, Filip**; Patent Department,
Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340
Beerse (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RQ, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*
- *of inventorship (Rule 4.17(iv)) for US only*

Published:

without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/20501 A2

BEST AVAILABLE COPY

(54) Title: POLYARYLCARBOXAMIDES USEFUL AS LIPID LOWERING AGENTS

(57) Abstract: Polyarylcaboxamide compounds of formula (I) are useful as lipid lowering agents.

POLYARYLCARBOXAMIDES USEFUL AS LIPID LOWERING AGENTS

The present invention is concerned with novel polyarylcaboxamides having
5 apolipoprotein B inhibiting activity and concomitant lipid lowering activity. The
invention further relates to methods for preparing such compounds,
pharmaceutical compositions comprising said compounds as well as the use of
said compounds as a medicine for the treatment of hyperlipidemia, obesity and
type II diabetes.

10

BACKGROUND OF THE INVENTION

Obesity is the cause of a myriad of serious health problems like the adult (48)
onset of diabetes and heart disease. In addition, the loss of weight is getting an
obsession among an increasing proportion of the human population.

15 The causal relationship between hypercholesterolemia, particularly that
associated with increased plasma concentrations of low density lipoproteins
(hereinafter referred as LDL) and very low density lipoproteins (hereinafter
referred as VLDL), and premature atherosclerosis and/or cardiovascular
disease is now widely recognized. However, a limited number of drugs are
20 presently available for the treatment of hyperlipidemia. Drugs primarily used for
the management of hyperlipidemia include bile acid sequestrant resins such as
cholestyramine and colestipol, fibric acid derivatives such as bezafibrate,
clofibrate, fenofibrate, ciprofibrate and gemfibrozil, nicotinic acid and cholesterol
synthesis inhibitors such as HMG Co-enzyme-A reductase inhibitors. The
25 inconvenience of administration (a granular form to be dispersed in water or
orange juice) and the major side-effects (gastro-intestinal discomfort and
constipation) of bile acid sequestrant resins constitute major drawbacks. Fibric
acid derivatives induce a moderate decrease (by 5 to 25%) of LDL cholesterol
(except in hypertriglyceridemic patients in whom initially low levels tend to
30 increase) and, although usually well tolerated, suffer from side-effects including
potentiation of warfarine, pruritus, fatigue, headache, insomnia, painful
reversible myopathy and stiffness in large muscle groups, impotency and
impaired renal function. Nicotinic acid is a potent lipid lowering agent resulting
in a 15 to 40% decrease in LDL cholesterol (and even 45 to 60% when
35 combined with a bile acid sequestrant resin) but with a high incidence of

troublesome side-effects related to the drug's associated vasodilatory action, such as headache, flushing, palpitations, tachychardia and occasional syncopes, as well as other side-effects such as gastro-intestinal discomfort, hyperucemia and impairment of glucose tolerance. Among the family of HMG

5 Co-enzyme-A reductase inhibitors, lovastatin and simvastatin are both inactive prodrugs containing a lactone ring which is hydrolyzed in the liver to form the corresponding active hydroxy-acid derivative. Inducing a reduction of LDL cholesterol by 35 to 45%, they are generally well tolerated with alow incidence of minor side effects. However there still remains a need for new lipid lowering

10 agents with improved efficiency and/or acting via other mechanisms than the above mentioned drugs.

Plasma lipoproteins are water-soluble complexes of high molecular weight formed from lipids (cholesterol, triglyceride, phospholipids) and apolipoproteins. Five major classes of lipoproteins that differ in the proportion of lipids and the

15 type of apolipoprotein, all having their origin in the liver and/or the intestine, have been defined according to their density (as measured by ultracentrifugation). They include LDL, VLDL, intermediate density lipoproteins (hereinafter referred as IDL), high density lipoproteins (hereinafter referred as HDL) and chylomicrons. Ten major human plasma apolipoproteins have been

20 identified. VLDL, which is secreted by the liver and contains apolipoprotein B (hereinafter referred as Apo-B), undergoes degradation to LDL which transports 60 to 70% of the total serum cholesterol. Apo-B is also the main protein component of LDL. Increased LDL-cholesterol in serum, due to oversynthesis or decreased metabolism, is causally related to atherosclerosis. In contrast high

25 density lipoproteins (hereinafter referred as HDL), which contain apolipoprotein A1, have a protective effect and are inversely correlated with the risk of a coronary heart disease. The HDL/LDL ratio is thus a convenient method of assessing the atherogenic potential of an individual's plasma lipid profile.

The two isoforms of apolipoprotein (apo) B, apo B-48 and apo B-100, are

30 important proteins in human lipoprotein metabolism. Apo B-48, so named because it appears to be about 48% the size of apo B-100 on sodium dodecyl sulfate-polyacrylamide gels, is synthesized by the intestine in humans. Apo B-48 is necessary for the assembly of chylomicrons and therefore has an obligatory role in the intestinal absorption of dietary fats. Apo B-100, which is

produced in the liver in humans, is required for the synthesis and secretion of VLDL. LDL, which contain about 2/3 of the cholesterol in human plasma, are metabolic products of VLDL. Apo B-100 is virtually the only protein component of LDL. Elevated concentrations of apo B-100 and LDL cholesterol in plasma
5 are recognized risk factors for developing atherosclerotic coronary artery disease.

A large number of genetic and acquired diseases can result in hyperlipidemia. They can be classified into primary and secondary hyperlipidemic states. The most common causes of the secondary
10 hyperlipidemias are diabetes mellitus, alcohol abuse, drugs, hypothyroidism, chronic renal failure, nephrotic syndrome, cholestasis and bulimia. Primary hyperlipidemias have also been classified into common hypercholesterolaemia, familial combined hyperlipidaemia, familial hypercholesterolaemia, remnant hyperlipidaemia, chylomicronaemia syndrome and familial hyper-
15 triglyceridaemia.

Microsomal triglyceride transfer protein (hereinafter referred as MTP) is known to catalyze the transport of triglyceride and cholesteryl ester by preference to phospholipids such as phosphatidylcholine. It was demonstrated by D.Sharp et al., *Nature* (1993) 365:65 that the defect causing
20 abetalipoproteinemia is in the MTP gene. This indicates that MTP is required for the synthesis of Apo B-containing lipoproteins such as VLDL, the precursor to LDL. It therefore follows that an MTP inhibitor would inhibit the synthesis of VLDL and LDL, thereby lowering levels of VLDL, LDL, cholesterol and triglyceride in humans. MTP inhibitors have been reported in Canadian patent
25 application No. 2,091,102 and in WO 96/26205. MTP inhibitors belonging to the class of polyarylcaboxamides have also been reported in U.S.Patent No. 5,760,246 as well as in WO-96/40640 and WO-98/27979.


One of the goals of the present invention is to provide an improved treatment for patients suffering from obesity or atherosclerosis, especially coronary
30 atherosclerosis and more generally from disorders which are related to atherosclerosis, such as ischaemic heart disease, peripheral vascular disease and cerebral vascular disease. Another goal of the present invention is to cause regression of atherosclerosis and inhibit its clinical consequences, particularly morbidity and mortality.

SUMMARY OF THE INVENTION

The present invention is based on the unexpected discovery that a class of novel polyarylcarboxamide compounds is acting as selective MTP inhibitors, i.e. is able to selectively block MTP at the level of the gut wall in mammals, and is therefore a promising candidate as a medicine, namely for the treatment of hyperlipidemia. The present invention additionally provides several methods for preparing such polyarylcarboxamide compounds, as well as pharmaceutical compositions including such compounds. Furthermore, the invention provides a certain number of novel compounds which are useful intermediates for the preparation of the therapeutically active polyarylcarboxamide compounds, as well as methods for preparing such intermediates. Finally, the invention provides a method of treatment of a condition selected from atherosclerosis, pancreatitis, obesity, hypercholesterolemia, hypertriglyceridemia, hyperlipidemia, diabetes and type II diabetes, comprising administering a therapeutically active polyarylcarboxamide compound to a mammal.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a family of novel polyarylcarboxamide compounds of formula (I)



the N-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

- Z_1 is selected from $(CH_2)_n$ wherein n is 1 to 3, CH_2CH_2O and OCH_2CH_2 ;
- Z_2 is $(CH_2)_m$ wherein m is 1 or 2;
- X_1 represents O, CH_2 , CO, NH, CH_2O , OCH_2 , CH_2S , SCH_2 or a direct bond;
- X_2 and X_3 are each independently selected from CH, N and a sp^2 carbon atom;
- R_1 is hydrogen or C_{1-4} alkyl;

- Ar¹ is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one or two R₃ substituents;
- 5 - Ar² is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one, two or three R₄ substituents;
- each R₂ and R₃ is independently selected from C₁₋₄alkyl, C₁₋₄alkyloxy, halo, trifluoromethyl;
- 10 - each R₄ is independently selected from C₁₋₄alkyl, C₁₋₄alkyloxy, halo, hydroxy, mercapto, cyano, nitro, C₁₋₄alkylthio or polyhaloC₁₋₆alkyl, amino, C₁₋₄alkylamino and di(C₁₋₄alkyl)amino;
- p¹ and p² are each 0 to 2;
- 15 - p³ is 0 to 3;
- X₁ and R₄ taken together with the aromatic rings Ar¹ and Ar² to which they are attached may form a fluoren-1-yl or a fluoren-4-yl group;
- A represents a C₁₋₆alkanediyl substituted with one or two groups selected from aryl, heteroaryl and C₃₋₁₀cycloalkyl; or when X₃ is CH, A may also
- 20 represent a nitrogen atom substituted with hydrogen, C₁₋₁₀alkyl, aryl, heteroaryl, arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl or C₃₋₁₀ cycloalkyl;
- B represents hydrogen; C₁₋₁₀alkyl; aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio,
- 25 C₁₋₁₀alkoxycarbonyl, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl; heteroaryl C₁₋₁₀alkyl; C₃₋₁₀cycloalkyl; polyhaloC₁₋₆alkyl; C₃₋₆ alkenyl; C₃₋₆ alkynyl; NR₆R₇; or OR₈;
- R₆ and R₇ each independently represent hydrogen, C₁₋₁₀alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo,
- 30 cyano, C₁₋₄ alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl, C₃₋₁₀cycloalkyl, C₇₋₁₀polycycloalkyl, polyhaloC₁₋₆alkyl, C₃₋₈alkenyl, C₃₋₈ alkynyl, fused benzo-C₅₋₈cycloalkyl, and

wherein R_6 and R_7 taken together with the nitrogen atom to which they are attached may form a C_{4-8} saturated heterocyclic radical; and

- R_8 represents C_{1-10} alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C_{1-4} alkyloxy, amino, C_{1-10} alkylamino, di(C_{1-10} alkyl)amino, C_{1-10} acyl, C_{1-10} alkylthio, C_{1-10} alkylaminocarbonyl and di(C_{1-10} alkyl)aminocarbonyl; aryl C_{1-10} alkyl; heteroaryl C_{1-10} alkyl; C_{3-10} cycloalkyl; C_{7-10} polycycloalkyl; polyhalo C_{1-6} alkyl; C_{3-8} alkenyl; C_{3-8} alkynyl; or fused benzo- C_{5-8} cycloalkyl.

Unless otherwise stated, as used in the foregoing definitions and hereinafter:

- 10 - halo is generic to fluoro, chloro, bromo and iodo;
- C_{1-4} alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, n-butyl, 1-methylethyl, 2-methylpropyl, 1,1-dimethylethyl and the like;
- C_{1-6} alkyl is meant to include C_{1-4} alkyl (as hereinabove defined) and the
15 higher homologues thereof having 5 or 6 carbon atoms, such as for instance 2-methylbutyl, n-pentyl, dimethylpropyl, n-hexyl, 2-methylpentyl, 3-methylpentyl and the like;
- C_{1-10} alkyl is meant to include C_{1-6} alkyl (as hereinabove defined) and the higher homologues thereof having 7 to 10 carbon atoms, such as for
20 instance heptyl, ethylhexyl, octyl, nonyl, decyl and the like;
- C_{3-10} cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl and cyclodecyl;
- polyhalo C_{1-6} alkyl is defined as polyhalosubstituted C_{1-6} alkyl, in particular C_{1-6} alkyl (as hereinabove defined) substituted with 2 to 13 halogen atoms
25 such as difluoromethyl, trifluoromethyl, trifluoroethyl, octafluoropentyl and the like;
- aryl is defined as mono- and polyaromatic groups such as phenyl or naphthalenyl optionally substituted with one to three substituents each independently selected from nitro, azido, cyano, halo, hydroxy, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{1-4} alkyloxy, polyhalo C_{1-6} alkyl, amino, mono- or
30 di(C_{1-6} alkyl)amino;
- heteroaryl is defined as mono- and polyheteroaromatic groups such as those including one or more heteroatoms selected from nitrogen, oxygen, sulfur and phosphorus, in particular pyridinyl, pyrazinyl, pyrimidinyl,

pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl, thienyl and the like, including all possible isomeric forms thereof;

- C₃₋₆alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 3 to 6 carbon atoms such as, for example, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl, 2-hexenyl and the like;
- C₃₋₆alkynyl defines straight and branched chain hydrocarbon radicals containing one triple bond and having from 3 to 6 carbon atoms such as, for example, 2-propynyl, 3-butylnyl, 2-butylnyl, 2-pentylnyl, 3-pentylnyl, 3-methyl-2-butylnyl, 3-hexynyl, 2-hexynyl and the like;
- C₄₋₈cycloalkenyl defines cyclic hydrocarbon radicals containing one double bond and having from 4 to 8 carbon atoms such as, for example cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl and the like;
- fused benzo-C₅₋₈cycloalkyl defines radicals such as, for instance, indanyl, 1,2,3,4-tetrahydronaphthalenyl, fluorenyl and the like;
- C₇₋₁₀ polycycloalkyl defines radicals having from 7 to 10 carbon atoms such as, for instance, norbornyl;
- C₁₋₆alkylamino defines primary amino radicals having from 1 to 6 carbon atoms such as, for example, methylamino, ethylamino, propylamino, isopropylamino, butylamino, isobutylamino and the like;
- di(C₁₋₆ alkyl)amino defines secondary amino radicals having from 1 to 6 carbon atoms such as, for example, dimethylamino, diethylamino, dipropylamino, di-isopropylamino, N-methyl-N'-ethylamino, N-ethyl-N'-propylamino and the like;
- C₁₋₆alkylthio defines a C₁₋₆alkyl group attached to a sulfur atom, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio and the like;
- C₁₋₆acyl defines a C₁₋₆alkyl group attached to a carbonyl group such as, for instance acetyl, propionyl, butyryl, isobutyryl and the like.

The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to include the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form and which may conveniently be obtained by treating the base form of such compounds with an

appropriate acid. Examples of such appropriate acids include, for instance, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 5 2-oxopropanoic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic (i.e. 2-hydroxybenzoic), p-aminosalicylic, pamoic and the like. Conversely the salt form can be converted by treatment with an appropriate 10 alkali into the free base form.

The term pharmaceutically acceptable addition salt as used hereinabove also includes the solvates which the compounds of formula (I) as well as their salts are able to form, such as for example hydrates, alcoholates and the like.

The N-oxide forms of the compounds of formula (I), are meant to include 15 those compounds wherein one or more nitrogen atoms are oxidized, using methods well known in the art for converting a trivalent nitrogen into its N-oxide form. Said N-oxidation reaction may usually be carried out by reacting the compound of formula (I) with 3-phenyl-2-(phenylsulfonyl)oxaziridine or with an appropriate organic or inorganic peroxide in at least one suitable solvent. 20 Appropriate inorganic peroxides include for example hydrogen peroxide and alkali metal or alkaline-earth metal peroxides, e.g. sodium or potassium peroxides. Appropriate organic peroxides may comprise peroxy acids such as, for example benzenecarboperoxoic acid or halo substituted benzenecarbo - peroxy acid (e.g. 3-chlorobenzene-carboperoxoic acid), peroxyalkanoic acids 25 (e.g. peroxyacetic acid) and alkylhydroperoxides (e.g. tert-butyl hydroperoxide). Suitable solvents for this reaction include for instance water, lower alcohols (e.g. ethanol and the like), hydrocarbons (e.g. toluene), ketones (e.g. 2-butanone), halogenated hydrocarbons (e.g. dichloromethane) and mixtures of such solvents.

30 The polyarylcaboxamide compounds of formula (I) may have at least one chiral center in the A group and/or the B group and/or the cyclic group including X_2 and X_3 .

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible isomeric forms which the compounds of formula (I) may possess.

Unless otherwise stated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. More particularly, stereogenic centers may have either the R- or S-
5 configuration; substituents on bivalent cyclic saturated radicals may have either the cis- or trans-configuration. The same definition applies to the various novel intermediates, as described herein, which are used to prepare the polyarylcarboxamide compounds of formula (I).

Pure stereoisomeric forms of the said compounds and intermediates are
10 defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure. In particular, the term "stereoisomerically pure" or "chirally pure" relates to compounds or intermediates having a stereoisomeric excess of at least 80% (i.e. at least 90% of one isomer and at most 10% of the other possible isomers), preferably at
15 least 90%, more preferably at least 94% and most preferably at least 97%. The terms "enantiomerically pure" and "diastereomerically pure" should be understood in a similar way, having regard to the enantiomeric excess, respectively the diastereomeric excess, of the mixture in question.

Consequently, if a mixture of enantiomers is obtained during any of the
20 following preparation methods, it can be separated by liquid chromatography using a suitable chiral stationary phase. Suitable chiral stationary phases are, for example, polysaccharides, in particular cellulose or amylose derivatives. Commercially available polysaccharide based chiral stationary phases are ChiralCel™ CA, OA, OB, OC, OD, OF, OG, OJ and OK, and Chiralpak™ AD,
25 AS, OP(+) and OT(+). Appropriate eluents or mobile phases for use in combination with said polysaccharide chiral stationary phases are hexane and the like, modified with an alcohol such as ethanol, isopropanol and the like.

The terms *cis* and *trans* are used herein in accordance with Chemical Abstracts nomenclature and refer to the position of the substituents on a ring
30 moiety.

The absolute stereochemical configuration of the polyarylcarboxamide compounds of formula (I) and of the intermediates used in their preparation may easily be determined by those skilled in the art while using well-known methods such as, for example, X-ray diffraction.

Furthermore, some polyarylcaboxamide compounds of formula (I) and some of the intermediates used in their preparation may exhibit polymorphism. It is to be understood that the present invention encompasses any polymorphic forms possessing properties useful in the treatment of the conditions noted
5 hereinabove.

A group of interesting compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply :

- a) R^1 is hydrogen;
 - 10 b) R^2 is hydrogen or C_{1-4} alkyl;
 - c) R^3 is hydrogen or C_{1-4} alkyl;
 - d) R^4 is hydrogen, C_{1-4} alkyl or trifluoromethyl;
 - e) p^1 is 1;
 - f) p^2 is 1;
 - 15 g) p^3 is 1;
 - h) the bivalent radical A is C_{1-6} alkanediyl substituted with one aryl group, in particular A is a methylene group substituted with phenyl;
 - i) B is C_{1-4} alkyloxy, or C_{1-10} alkylamino.
- 20 Interesting compounds are those compounds of formula (I) wherein Z_1 , Z_2 , X_2 and X_3 taken together form a six-membered heterocycle.

Particular compounds are those compounds of formula (I) wherein radical B represents methyloxy or ethyloxy.















25 Other particular compounds are those compounds of formula (I) wherein R_2 and/or R_3 are/is C_{1-4} alkyl.


Yet other particular compounds are those compounds of formula (I) wherein R_4
30 is C_{1-4} alkyl or trifluoromethoxy.

More preferred compounds are those particular compounds of formula (I) wherein Z_1 , Z_2 , X_2 and X_3 taken together form a piperidine or piperazine group and X_1 is a direct bond.

35 More preferred compounds of formula (I) are those compounds wherein R_2 and R_3 are each hydrogen and R_4 is hydrogen, trifluoromethyl, chloro or tert-butyl.

Most preferred compound of formula (I) are






and the stereoisomeric forms, the pharmaceutically acceptable acid addition salts, or the *N*-oxides thereof.


One advantage of the present invention is the easiness with which the compounds of formula (I) can be manufactured by a high number of different processes. Some of these processes will now be described in details, without
 5 pretending to provide an exhaustive list of the methods for preparing the said compounds.

A first process for preparing a polyarylcaboxamide compound according to this invention is a process wherein an intermediate phenylene amine having the
 10 formula



(II)

wherein Z_1 , Z_2 , X_2 , X_3 , p^1 , R_1 , R_2 , A and B are as defined in formula (I), is
 15 reacted with a polyarylcaboxylic acid or halide having the formula (III),




(III)

wherein X_1 , Ar^1 , Ar^2 , p^2 , p^3 , R_3 and R_4 are as defined in formula (I) and Y_1 is
 20 selected from hydroxy and halo, in at least one reaction-inert solvent and optionally in the presence of a suitable base, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. In case Y_1 is
 25 (III) by adding an effective amount of a reaction promoter. Non-limiting examples of such reaction promoters include carbonyldiimidazole, diimides

such as N,N'-dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and functional derivatives thereof. For this type of acylation procedure, it is preferred to use a polar aprotic solvent such as, for instance, methylene chloride. Suitable bases for carrying out this first process include

5 tertiary amines such as triethylamine, triisopropylamine and the like. Suitable temperatures for carrying out the first process of the invention typically range from about 20°C to about 140°C, depending on the particular solvent used, and will most often be the boiling temperature of the said solvent.


- 10 A second process for preparing a polyarylcaboxamide compound according to this invention, wherein X_3 is nitrogen, is a process wherein an intermediate having the formula



(IV)

15

wherein Z_1 , Z_2 , X_1 , X_2 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I) and X_3 is nitrogen, is reacted with a reactant having the formula (V)




(V)

- 20 wherein A and B are as defined in formula (I) and Y_2 is selected from halo, tosyloxy, mesyloxy, naphthylsulfonyloxy, or $-A-Y_2$ is R'_5COR_5 , wherein R_5 and R'_5 are such that the radical R'_5CHR_5 is encompassed by the definition of A in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one suitable nucleophilic substitution activator and/or a suitable
- 25 base, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. When Y_2 is halo, the alkylation coupling procedure may for instance be effected in the presence of sodium or potassium carbonate or a tertiary amine such as triethylamine or diisopropylethylamine in a solvent such

as dimethylformamide or methylisopropylketone and optionally in the presence of a catalytic amount of potassium iodide in order to enhance nucleophilic substitution. Intermediates of formula (IV) wherein X_3 is nitrogen may also be reductively N-alkylated by means of an aldehyde or a ketone of formula (V) wherein -A-Y₂ is R'₅COR₅, thus forming compounds of formula (I). Said reductive N-alkylation may be performed in a reaction-inert solvent such as for example toluene, methanol, tetrahydrofuran or a mixture thereof, and in the presence of a reducing agent. Non-limiting examples of such reducing agents include hydrogen and borohydrides, e.g. sodium borohydride, zinc borohydride, lithium borohydride, sodium cyanoborohydride, triacetoxo borohydride and the like. When a borohydride is used as the reducing agent, it may be convenient to perform the N-alkylation in the additional presence of a catalyst. Non-limiting examples of such catalysts include transition metal alkoxides, e.g. titanium(IV)isopropoxide, titanium(IV)-n-butoxide and the like, as disclosed in *J.Org.Chem.* (1990), 55:2552-4. When hydrogen is used as the reducing agent, it may be convenient to perform the N-alkylation in the additional presence of a catalyst. Non-limiting examples of catalysts suitable for this purpose include a noble metal supported on a carrier such as for instance palladium-on-charcoal or platinum-on-charcoal. The formation of a Schiff base in a first step of the said reductive N-alkylation reaction may be further enhanced by the additional presence of a suitable reagent such as aluminium terbutoxide, calcium oxide, calcium hydride and the like. An appropriate catalyst-poison, e.g. thiophene, butanethiol, quinoline sulfur or the like, may also be added to the reaction mixture in order to prevent undesired hydrogenation of certain functional groups in the reactants and/or the reaction product. Stirring and optionally elevated temperature and/or pressure may further enhance the rate of such a reaction.

A third process for preparing a polyarylcarboxamide compound of the invention is a process wherein an intermediate having the formula




(VI)

wherein Z_1 , Z_2 , X_1 , X_2 , X_3 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 , R_4 and A are as defined in formula (I) and Y_3 is selected from halo and hydroxy, is reacted with a reactant of the formula BH, wherein B is NR_6R_7 or OR_8 and R_6 , R_7 and R_8 are as defined in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one suitable coupling reagent and/or a suitable base,, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. In case Y_3 is hydroxy, it may be convenient to activate the carboxylic acid of formula (VI) by adding an effective amount of a reaction promoter. Non-limiting examples of such reaction promoters include carbonyldiimidazole, diimides such as N,N'-dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbo-diimide, and functional derivatives thereof. In case a chirally pure reactant of formula (V) is used, a fast and enantiomerization-free reaction of the intermediate of formula (VI) with the said reactant may be performed in the further presence of an effective amount of a compound such as hydroxybenzotriazole, benzotriazolyloxytris(dimethylamino)phosphonium hexafluorophosphate, tetrapyrrolidinophosphonium hexafluorophosphate, bromotripyrrolidinophosphonium hexafluorophosphate, or a functional derivative thereof, such as disclosed by D. Hudson, *J.Org.Chem.* (1988), 53:617. In case Y_3 is hydroxy and B is OR_8 , then the esterification reaction may conveniently be performed in the presence of an effective amount of an acid such as sulfuric acid and the like.


25 A fourth process for preparing a polyarylcaboxamide compound according to this invention wherein X_3 is nitrogen and wherein A is a group suitable for a Michael addition reaction, is a process wherein an intermediate having the

formula (IV), wherein X_3 is nitrogen, is reacted with a reactant of the formula (VII)




(VII)


- 5 wherein B is as defined in formula (I), and Y_4 and Y_4' are such that the radical



- is encompassed by the definition of A in formula (I), in at least one reaction-inert solvent, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing
- 10 stereochemically isomeric forms thereof. The reaction-inert solvent may be, for instance, dimethylformamide or methanol and the reaction temperature may be the boiling point of the said solvent. Contrary to most of the alternative processes for preparing the polyarylcaboxamide compounds of this invention, this fourth process does not require the presence of a catalyst or otherwise
- 15 coupling reagent in order to quantitatively yield the target compound. A base such as sodium carbonate, potassium carbonate, cesium carbonate and the like may optionally be added to the reaction mixture. As conventionally defined in the art, A preferably is an α,β -unsaturated carbonyl compound, such as a ketone or an ester, the β carbon atom of which is susceptible of a nucleophilic
- 20 attack. Non-limiting examples of groups A suitable for such a Michael addition reaction include for instance:




- A fifth process for preparing a polyarylcaboxamide compound according to this
- 25 invention is a process wherein an intermediate having the formula (VIII)



(VIII)


wherein X_1 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I) and Y_5 is selected from halo, $B(OH)_2$, alkylboronates and cyclic analogues thereof, is reacted with a reactant having the formula (IX)




(IX)

wherein Z_1 , Z_2 , X_3 , A and B are as defined in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one transition metal coupling reagent and/or at least one suitable ligand, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. This type of reaction being known in the art as the Buchwaldt reaction, reference to the applicable metal coupling reagents and/or suitable ligands, e.g. palladium compounds such as palladium tetra(triphenylphosphine), tris(dibenzylidene-acetone dipalladium, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and the like, may be found for instance in *Tetrahedron Letters* (1996) 37(40) 7181-7184 and *J.Am.Chem.Soc.* (1996) 118:7216. If Y_5 is $B(OH)_2$, an alkylboronate or a cyclic analogue thereof, then cupric acetate should be used as the coupling reagent, according to *Tetrahedron Letters* (1998) 39:2933-6.

A sixth process for preparing a polyarylcaboxamide compound according to this invention is a process wherein an intermediate having the formula (X)




wherein p^1 , p^2 , p^3 , Ar^1 , Ar^2 , X_1 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I),
is reacted with a reactant having the formula (XI)




wherein Z_1 , Z_2 , X_3 , A and B are as defined in formula (I), one of Y_6 and Y_7 is selected from bromo, iodo and trifluoromethylsulfonate and the other of Y_6 and Y_7 is selected from tri(C_{1-4} alkyl) tin, $B(OH)_2$, alkylboronates and cyclic analogues thereof, in at least one reaction-inert solvent and optionally in the presence of at least one transition metal coupling reagent and/or at least one suitable ligand such as palladium associated with triphenylphosphine, triphenylarsine and the like, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. This type of reaction being known in the art as the Stille reaction or the Suzuki reaction, reference to the applicable transition metal coupling reagents and/or suitable ligands may be found for instance in *Syn.Letters* (1998)6,671-5, in *Chem.Rev.*(1999)99(6)1549-1581 and in *The Stille Reaction* (John Wiley & Sons, Inc.) ISBN 0-471-31273-8.

A seventh process for preparing a polyarylcaboxamide compound according to this invention is a process wherein an intermediate having the formula (XII)




(XII)

wherein Z_1 , Z_2 , X_2 , X_3 , p^1 , p^2 , Ar^1 , R_1 , R_2 , R_3 , A and B are as defined in formula
 5 (I) and Y_8 is selected from bromo, iodo and trifluoromethylsulfonate, is reacted
 either with an aryl-boric acid having the formula (XIII a)



(XIII-a)

wherein p^3 , Ar^2 and R_4 are as defined in formula (I), or with an aryl-tin reactant
 10 having the formula (XIII b)




(XIII-b)

wherein R_3 , R_4 , Ar^2 and p^3 are as defined in formula (I), in at least one reaction-
 inert solvent and optionally in the presence of at least one transition metal
 15 coupling reagent and/or at least one suitable ligand,, the said process further
 optionally comprising converting a compound of formula (I) into an addition salt
 thereof, and/or preparing or stereochemically isomeric forms thereof. This type
 of reaction being known in the art as the Stille reaction or the Suzuki reaction,
 reference to the applicable transition metal coupling reagents and/or suitable
 20 ligands may be found for instance in the literature cited above. Exemplary
 coupling reagents for this process are e.g. bis(triphenylphospine)
 dichloropalladium and diacetylpalladium. Exemplary reaction-inert solvents for
 this reaction are 1,4-dioxane, toluene, dimethylformamide, tetrahydrofuran,
 dimethylether and the like.


Furthermore, a process for preparing a polyarylcaboxamide compound of the formula (I) wherein B is NR_6R_7 , from a polyarylcaboxamide compound of the formula (I) wherein B is OR_5 is a process comprising in a first step hydrolyzing the latter and in a second step reacting the resulting corresponding carboxylic acid with an amine having the formula HNR_6R_7 in at least one reaction-inert solvent and further optionally comprising converting the resulting compound of formula (I) wherein B is NR_6R_7 into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof. The hydrolysis first step is preferably effected in an acidic medium such as strongly concentrated hydrochloric acid and optionally in the presence of an organic solvent such as dioxane.

The compounds of formula (I) can also conveniently be prepared using solid phase synthesis techniques. In general, solid phase synthesis involves reacting an intermediate in a synthesis with a polymer support. This polymer supported intermediate can then be carried on through a number of synthetic steps. After each step, impurities are removed by filtering the resin and washing it numerous times with various solvents. At each step the resin can be split up to react with various intermediates in the next step thus allowing for the synthesis of a large number of compounds. After the last step in the procedure the resin is treated with a reagent or process to cleave the resin from the sample. More detailed explanation of the techniques used in solid phase chemistry are described in for example "The Combinatorial Index" (B.Bunin, Academic Press) and Novabiochem's 1999 Catalogue & Peptide Synthesis Handbook (Novabiochem AG, Switzerland) both incorporated herein by reference.


Furthermore, the invention provides compounds of any of the formulae (II), (III), (IV), (VI), (VIII), (X) and (XII) hereunder:




(11)




(III)




(IV)




(VI)



(VIII)



(X)



wherein :

- Z_1 is selected from $(CH_2)_n$ wherein n is 1 to 3, CH_2CH_2O and OCH_2CH_2 ;
- 5 - Z_2 is $(CH_2)_m$ wherein m is 1 or 2;
- X_1 represents O, CH_2 , CO, NH, CH_2O , OCH_2 , CH_2S , SCH_2 or a direct bond;
- X_2 and X_3 are each independently selected from CH, N and a sp^2 carbon atom;
- R_1 is hydrogen or C_{1-4} alkyl;
- 10 - Ar^1 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one or two R_3 substituents;
- Ar^2 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one, two or three R_4 substituents;
- 15 - each R_2 and R_3 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo and trifluoromethyl;
- 20 - each R_4 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo, hydroxy, mercapto, cyano, nitro, C_{1-4} alkylthio or polyhalo C_{1-6} alkyl, amino, C_{1-4} alkylamino and di(C_{1-4} alkyl)amino;
- p^1 and p^2 are each 0 to 2;
- p^3 is 0 to 3;
- 25 - X_1 and R_4 taken together with the aromatic rings to which they are attached may form a fluoren-1-yl or a fluoren-4-yl group;
- A represents a C_{1-6} alkanediyl optionally substituted with one or two groups selected from aryl, heteroaryl and C_{3-10} cycloalkyl; oxygen; or a direct bond;

- B represents hydrogen; C₁₋₁₀alkyl; aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkoxycarbonyl, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl; heteroarylC₁₋₁₀alkyl; C₃₋₁₀ cycloalkyl; polyhaloC₁₋₆alkyl; C₃₋₆ alkenyl; C₃₋₆ alkynyl; NR₆R₇; or OR₈;
- R₆ and R₇ each independently represent hydrogen, C₁₋₁₀alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl, C₃₋₁₀cycloalkyl, C₇₋₁₀polycycloalkyl, polyhaloC₁₋₆alkyl, C₃₋₈alkenyl, C₃₋₈alkynyl, fused benzo-C₅₋₈cycloalkyl, and wherein R₆ and R₇ taken together with the nitrogen atom to which they are attached may form a C₄₋₈ saturated heterocyclic radical;
- R₈ represents C₁₋₁₀alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl; heteroarylC₁₋₁₀alkyl; C₃₋₁₀cycloalkyl; C₇₋₁₀ polycycloalkyl; polyhaloC₁₋₆alkyl; C₃₋₈alkenyl; C₃₋₈alkynyl; or fused benzo-C₅₋₈ cycloalkyl;
- when X₃ is CH, A may also represent a nitrogen atom substituted with hydrogen, C₁₋₁₀alkyl, aryl, heteroaryl, arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl or C₃₋₁₀cycloalkyl;
- Y₁ and Y₃ are each independently selected from hydroxy and halo;
- Y₅ is selected from halo, B(OH)₂, alkylboronates and cyclic analogues thereof; and
- Y₆ and Y₈ are each independently selected from bromo, iodo and trifluoromethylsulfonate,

which are useful as intermediates for the preparation of the polyarylcarboxamide compounds of the present invention. In turn, the present invention provides methods for preparing the above-mentioned families of intermediate compounds, such as disclosed in the foregoing examples.

The polyarylcarboxamide compounds of formula (I), the *N*-oxide forms, the pharmaceutically acceptable salts and stereoisomeric forms thereof possess

favourable apolipoprotein B inhibiting activity and concomitant lipid lowering activity. Therefore the present compounds are useful as a medicine especially in a method of treating patients suffering from hyperlipidemia, obesity, atherosclerosis or type II diabetes. In particular the present compounds may
5 be used for the manufacture of a medicine for treating disorders caused by an excess of very low density lipoproteins (VLDL) or low density lipoproteins (LDL), and especially disorders caused by the cholesterol associated with said VLDL and LDL.

10 The causal relationship between hypercholesterolemia – particularly that associated with increased plasma concentrations of low density lipoproteins (LDL) and very low density lipoproteins (VLDL) – and premature atherosclerosis and cardiovascular disease is well established. VLDL is secreted by the liver and contains apolipoprotein B (apo-B); these particles undergo degradation in
15 the circulation to LDL, which transports about 60 to 70% of the total serum cholesterol. Apo-B is also the principal protein component of LDL. Increased LDL-cholesterol in serum, due to oversynthesis or decreased metabolism, is causally related to atherosclerosis. In contrast, high density lipoproteins (HDL) which contain apolipoprotein A1, have a protective effect and are inversely
20 correlated with risk of coronary heart disease. The HDL/LDL ratio is thus a convenient method of assessing the atherogenic potential of an individual's plasma lipid profile.

The principal mechanism of action of the compounds of formula (I) appears to
25 involve inhibition of MTP (microsomal triglyceride transfer protein) activity in hepatocytes and intestinal epithelial cells, resulting in decreased VLDL and chylomicron production, respectively. This is a novel and innovative approach to hyperlipidemia, and is expected to lower LDL-cholesterol and triglycerides through reduced hepatic production of VLDL and intestinal production of
30 chylomicrons.

A large number of genetic and acquired diseases can result in hyperlipidemia. They can be classified into primary and secondary hyperlipidemic states. The most common causes of the secondary hyperlipidemias are diabetes mellitus,
35 alcohol abuse, drugs, hypothyroidism, chronic renal failure, nephrotic

syndrome, cholestasis and bulimia. Primary hyperlipidemias are common hypercholesterolaemia, familial combined hyperlipidaemia, familial hypercholesterolaemia, remnant hyperlipidaemia, chylomicronaemia syndrome, familial hypertriglyceridaemia. The present compounds may also be used to
5 prevent or treat patients suffering from obesitas or from atherosclerosis, especially coronary atherosclerosis and more in general disorders which are related to atherosclerosis, such as ischaemic heart disease, peripheral vascular disease, cerebral vascular disease. The present compounds may cause regression of atherosclerosis and inhibit the clinical consequences of
10 atherosclerosis, particularly morbidity and mortality.

In view of the utility of the compounds of formula (I), it follows that the present invention also provides a method of treating warm-blooded animals, including humans, (generally called herein patients) suffering from disorders caused by
15 an excess of very low density lipoproteins (VLDL) or low density lipoproteins (LDL), and especially disorders caused by the cholesterol associated with said VLDL and LDL. Consequently a method of treatment is provided for relieving patients suffering from conditions, such as, for example, hyperlipidemia, obesity, atherosclerosis or type II diabetes.

20 Apo B-48, synthesized by the intestine, is necessary for the assembly of chylomicrons and therefore has an obligatory role in the intestinal absorption of dietary fats. The present invention provides polyarylcarboxamide compounds which are acting as selective MTP inhibitors at the level of the gut wall.

25 Additionally the present invention provides pharmaceutical compositions comprising at least one pharmaceutically acceptable carrier and a therapeutically effective amount of a polyarylcarboxamide compound having the formula (I).

30 In order to prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, in base or addition salt form, as the active ingredient is combined in intimate admixture with at least one pharmaceutically acceptable carrier, which carrier may take a wide variety of
35 forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable,

preferably, for oral administration, rectal administration, percutaneous administration or parenteral injection.

For example in preparing the compositions in oral dosage form, any of the usual liquid pharmaceutical carriers may be employed, such as for instance
5 water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid pharmaceutical carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their easy administration, tablets and capsules represent the most
10 advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral injection compositions, the pharmaceutical carrier will mainly comprise sterile water, although other ingredients may be included in order to improve solubility of the active ingredient. Injectable solutions may be prepared for instance by using a
15 pharmaceutical carrier comprising a saline solution, a glucose solution or a mixture of both. Injectable suspensions may also be prepared by using appropriate liquid carriers, suspending agents and the like. In compositions suitable for percutaneous administration, the pharmaceutical carrier may optionally comprise a penetration enhancing agent and/or a suitable wetting
20 agent, optionally combined with minor proportions of suitable additives which do not cause a significant deleterious effect to the skin. Said additives may be selected in order to facilitate administration of the active ingredient to the skin and/or be helpful for preparing the desired compositions. These topical compositions may be administered in various ways, e.g., as a transdermal
25 patch, a spot-on or an ointment. Addition salts of the compounds of formula (I), due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the pharmaceutical compositions of the invention in dosage unit form for ease of administration and uniformity of
30 dosage. "Dosage unit form" as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined amount of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills,

powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

For oral administration, the pharmaceutical compositions of the present invention may take the form of solid dose forms, for example, tablets (both
5 swallowable and chewable forms), capsules or gelcaps, prepared by conventional means with pharmaceutically acceptable excipients and carriers such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone, hydroxypropylmethylcellulose and the like), fillers (e.g. lactose, microcrystalline cellulose, calcium phosphate and the like), lubricants (e.g. magnesium stearate,
10 talc, silica and the like), disintegrating agents (e.g. potato starch, sodium starch glycollate and the like), wetting agents (e.g. sodium laurylsulphate) and the like. Such tablets may also be coated by methods well known in the art.

Liquid preparations for oral administration may take the form of e.g. solutions, syrups or suspensions, or they may be formulated as a dry product
15 for admixture with water and/or another suitable liquid carrier before use. Such liquid preparations may be prepared by conventional means, optionally with other pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, methylcellulose, hydroxypropylmethylcellulose or hydrogenated edible fats), emulsifying agents (e.g. lecithin or acacia), non-aqueous carriers
20 (e.g. almond oil, oily esters or ethyl alcohol), sweeteners, flavours, masking agents and preservatives (e.g. methyl or propyl p-hydroxybenzoates or sorbic acid).

Pharmaceutically acceptable sweeteners useful in the pharmaceutical compositions of the invention comprise preferably at least one intense
25 sweetener such as aspartame, acesulfame potassium, sodium cyclamate, alitame, a dihydrochalcone sweetener, monellin, stevioside sucralose (4,1',6'-trichloro-4,1',6'-trideoxygalactosucrose) or, preferably, saccharin, sodium or calcium saccharin, and optionally at least one bulk sweetener such as sorbitol, mannitol, fructose, sucrose, maltose, isomalt, glucose, hydrogenated glucose
30 syrup, xylitol, caramel or honey. Intense sweeteners are conveniently used in low concentrations. For example, in the case of sodium saccharin, the said concentration may range from about 0.04% to 0.1% (weight/volume) of the final formulation. The bulk sweetener can effectively be used in larger

concentrations ranging from about 10% to about 35%, preferably from about 10% to 15% (weight/volume).

The pharmaceutically acceptable flavours which can mask the bitter tasting ingredients in the low-dosage formulations are preferably fruit flavours such as
5 cherry, raspberry, black currant or strawberry flavour. A combination of two flavours may yield very good results. In the high-dosage formulations, stronger pharmaceutically acceptable flavours may be required such as Caramel Chocolate, Mint Cool, Fantasy and the like. Each flavour may be present in the final composition in a concentration ranging from about 0.05% to 1%
10 (weight/volume). Combinations of said strong flavours are advantageously used. Preferably a flavour is used that does not undergo any change or loss of taste and/or color under the circumstances of the formulation.

The polyarylcarboxamide compounds of this invention may be formulated for parenteral administration by injection, conveniently intravenous, intra-muscular
15 or subcutaneous injection, for example by bolus injection or continuous intravenous infusion. Formulations for injection may be presented in unit dosage form, e.g. in ampoules or multi-dose containers, including an added preservative. They may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such
20 as isotoning, suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be present in powder form for mixing with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The polyarylcarboxamide compounds of this invention may also be formulated in rectal compositions such as suppositories or retention enemas,
25 e.g. containing conventional suppository bases such as cocoa butter and/or other glycerides.

The polyarylcarboxamide compounds of this invention may be used in conjunction with other pharmaceutical agents, in particular the pharmaceutical compositions of the present invention may further comprise at least one
30 additional lipid-lowering agent, thus leading to a so-called combination lipid-lowering therapy. The said additional lipid-lowering agent may be, for instance, a known drug conventionally used for the management of hyperlipidaemia such as e.g. a bile acid sequestrant resin, a fibric acid derivative or nicotinic acid as previously mentioned in the background of the invention. Suitable additional

lipid-lowering agents also include other cholesterol biosynthesis inhibitors and cholesterol absorption inhibitors, especially HMG-CoA reductase inhibitors and HMG-CoA synthase inhibitors, HMG-CoA reductase gene expression inhibitors, CETP inhibitors, ACAT inhibitors, squalene synthetase inhibitors and the like.

5 Any HMG-CoA reductase inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term "HMG-CoA reductase inhibitor" as used herein, unless otherwise stated, refers to a compound which inhibits the biotransformation of hydroxymethylglutaryl-coenzyme A to mevalonic acid as catalyzed by the enzyme HMG-CoA
10 reductase. Such inhibition may be determined readily by one skilled in the art according to standard assays, i.e. Methods of Enzymology (1981) 71:455-509. Exemplary compounds are described e.g. in U.S.Patent No.4,231,938 (including lovastatin), U.S.Patent No. 4,444,784 (including simvastatin), U.S.Patent No. 4,739,073 (including fluvastatin), U.S.Patent No. 4,346,227
15 (including pravastatin), EP-A-491,226 (including rivastatin) and U.S.Patent No. 4,647,576 (including atorvastatin).

Any HMG-CoA synthase inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term "HMG-CoA synthase inhibitor" as used herein, unless otherwise stated, refers to a compound which
20 inhibits the biosynthesis of hydroxymethylglutaryl-coenzyme A from acetyl-coenzyme A and acetoacetyl-coenzyme A, catalyzed by the enzyme HMG-CoA synthase. Such inhibition may be determined readily by one skilled in the art according to standard assays, i.e. Methods of Enzymology (1985) 110:19-26. Exemplary compounds are described e.g. in U.S.Patent No. 5,120,729 relating
25 to beta-lactam derivatives, U.S.Patent No. 5,064,856 relating to spiro-lactone derivatives and U.S.Patent No. 4,847,271 relating to oxetane compounds.

Any HMG-CoA reductase gene expression inhibitor may be used as the second compound in the combination therapy aspect of this invention. These agents may be HMG-CoA reductase transcription inhibitors that block the
30 transcription of DNA or translation inhibitors that prevent translation of mRNA coding for HMG-CoA reductase into protein. Such inhibitors may either affect transcription or translation directly or may be biotransformed into compounds having the above-mentioned attributes by one or more enzymes in the cholesterol biosynthetic cascade or may lead to accumulation of a metabolite

having the above-mentioned activities. Such regulation may be determined readily by one skilled in the art according to standard assays, i.e. Methods of Enzymology (1985) 110:9-19. Exemplary compounds are described e.g. in U.S. Patent No. 5,041,432 and E.I. Mercer, *Prog. Lip. Res.* (1993) 32:357-416.

5 Any CETP inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term "CETP inhibitor" as used herein, unless otherwise stated, refers to a compound which inhibits the cholesteryl ester transfer protein (CETP) mediated transport of various cholesteryl esters and triglycerides from HDL to LDL and VLDL. Exemplary
10 compounds are described e.g. in U.S. Patent No. 5,512,548, in *J. Antibiot.* (1996) 49(8):815-816 and *Bioorg. Med. Chem. Lett.* (1996) 6:1951-1954.

Any ACAT inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term "ACAT inhibitor" as used
15 herein, unless otherwise stated, refers to a compound which inhibits the intracellular esterification of dietary cholesterol by the enzyme acyl CoA:cholesterol acyltransferase. Such inhibition may be determined readily by one skilled in the art according to standard assays, i.e. the method of Heider et al., *Journal of Lipid Research* (1983) 24:1127. Exemplary compounds are
20 described e.g. in U.S. Patent No. 5,510,379, in WO 96/26948 and WO 96/10559.

Any squalene synthetase inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term "squalene synthetase inhibitor" as used herein, unless otherwise stated, refers to a
25 compound which inhibits the condensation of two molecules of farnesylpyrophosphate to form squalene, catalyzed by the enzyme squalene synthetase. Such inhibition may be determined readily by one skilled in the art according to standard methods, i.e. Methods of Enzymology (1985) 110:359-373. Exemplary compounds are described e.g. in EP-A-567,026, in EP-A-
30 645,378 and in EP-A- 645,377.

Those of skill in the treatment of hyperlipidemia will easily determine the therapeutically effective amount of a polyarylcarboxamide compound of this invention from the test results presented hereinafter. In general it is contemplated that a therapeutically effective dose will be from about 0.001

mg/kg to about 5 mg/kg of body weight, more preferably from about 0.01 mg/kg to about 0.5 mg/kg of body weight of the patient to be treated. It may be appropriate to administer the therapeutically effective dose in the form of two or more sub-doses at appropriate intervals throughout the day. Said sub-doses
5 may be formulated as unit dosage forms, for example each containing from about 0.1 mg to about 350 mg, more particularly from about 1 to about 200 mg, of the active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular polyarylcaboxamide compound of formula (I) used, the particular condition
10 being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as the other medication (including the above-mentioned additional lipid-lowering agents), the patient may be taking, as is well known to those skilled in the art. Furthermore, said effective daily amount may be lowered or increased depending on the
15 response of the treated patient and/or depending on the evaluation of the physician prescribing the polyarylcaboxamide compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.

20 Experimental part

In the procedures described hereinafter the following abbreviations were used: "ACN" stands for acetonitrile; "THF" stands for tetrahydrofuran; "DCM" stands for dichloromethane; "DIPE" stands for diisopropylether; "DMF" means *N,N*-dimethyl-formamide; "NMP" means *N*-methyl-2-pyrrolidone; "TFA" means
25 trifluoroacetic acid; "TIS" means triisopropylsilane; "DIPEA" means diisopropylethylamine; "TMSOTf" means trimethylsilyl triflate and "MIK" means methyl isobutyl ketone. Extrelut™ is a product of Merck KgaA, Darmstadt, Germany, and is a short column comprising diatomaceous earth.

30 A. Preparation of intermediate compounds

EXAMPLE A.1

α -Bromo-methyl ester benzeneacetic acid (0.026 mole) was added dropwise to a mixture of 1-(4-nitrophenyl)piperazine (0.028 mole) and Na₂CO₃ (0.024 mole) in DMF (150 ml) while stirring. The mixture was stirred at room temperature for
35 66 hours, poured out into ice water (500 ml) and stirred for 30 minutes. The precipitate was filtered off and dissolved in DCM. The organic solution was

dried, filtered and the solvent was evaporated. This fraction was purified over silica gel on a glass filter (eluent: CH₂Cl₂/CH₃OH 99/1). The pure fractions were collected and the solvent was evaporated. Part of this fraction was stirred in ethanol. The precipitate was filtered off and dried, yielding 0.04 g of (±)-methyl
5 4-(4-nitrophenyl)-α-phenyl-1-piperazineacetate (intermediate 1, melting point 92°C).

EXAMPLE A.2

A mixture of intermediate (1) (0.026 mole) and KOH (0.13 mole) in ethanol
10 (150 ml) was stirred at room temperature for 18 hours, heated at 50°C for 2.5 hours and cooled to room temperature. The precipitate was filtered off, stirred in 2-propanol, filtered off, washed three times with 2-propanol and dried. This fraction was stirred and refluxed in 2-propanol. HCl/2-propanol 6N (19.94 ml) was added. The mixture was stirred and refluxed, filtered warm and stirred in
15 water (350 ml). The precipitate was filtered off and dried, yielding 5.94 g of (±)-4-(4-nitrophenyl)-α-phenyl-1-piperazineacetic acid monohydrate (intermediate 2).

EXAMPLE A.3

A mixture of intermediate (1) (0.0136 mole) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.02 mole) in DCM (125 ml) was stirred for two hours to give mixture (1). A mixture of 2,2,2-trifluoroethylamine (0.014 mole) in DCM (25 ml) was stirred. Triethylamine (1.5 g) was added and the mixture was stirred for 5 minutes to give mixture (2). Mixtures (1) and (2) were combined. The resulting
25 mixture was stirred overnight and washed with water. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected and the solvent was evaporated. This fraction was purified by column chromatography over silica gel (eluent:
30 DCM/hexane/ethyl acetate 50/20/30). The pure fractions were collected and the solvent was evaporated, yielding 2.3g (±)-4-(4-nitrophenyl)-α-phenyl-N-(2,2,2-trifluoroethyl)-1-piperazineacetamide (intermediate 3).

EXAMPLE A.4

A mixture of intermediate (3) (0.0054 mole) in methanol (150 ml) was hydrogenated with Pd/C 10% (1 g) as a catalyst in the presence of a 4% thiophene solution (1 ml). After uptake of hydrogen (2 equivalents), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 1.5 g of (±)-4-(4-

aminophenyl)- α -phenyl-*N*-(2,2,2-trifluoroethyl)-1-piperazineacetamide
(intermediate 4, melting point 136°C).

EXAMPLE A.5

- 5 A mixture of 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carbonyl chloride (0.185 mole) in DCM (1500 ml) and triethylamine (50 ml) was stirred on an ice-bath for 5 minutes. 4-[4-(Phenylmethyl)-1-piperazinyl]-benzenamine (0.37 mole) in DCM (500 ml) was added dropwise. The mixture was stirred for 3 hours. The organic layer was separated, washed with water, dried, filtered and the solvent was
10 evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 99.8 g of *N*-[4-[4-(phenylmethyl)-1-piperazinyl]phenyl]-4'-(trifluoro-methyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 5, melting point 180°C).

EXAMPLE A.6

- 15 A mixture of intermediate (5) (0.19 mole) in methanol (600 ml) and THF (600 ml) was hydrogenated overnight with Pd/C 10% (3 g) as a catalyst. After uptake of hydrogen (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in DIPE. The precipitate was
20 filtered off, dried, and dissolved in water. The mixture was alkalized with Na₂CO₃ and then extracted with DCM. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 40 g of *N*-[4-(1-piperazinyl)phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide
25 (intermediate 6).

EXAMPLE A.7

- A mixture of 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.09 mole) in DCM (500 ml) and DMF (5 ml) was stirred. Ethanedioyl dichloride (0.09 mole)
30 was added dropwise. The mixture was stirred for 1 hour to give mixture (1). A mixture of 4-[1-(phenylmethyl)-4-piperidinyl]-benzenamine (0.046 mole) in DCM (500 ml) and triethylamine (20 ml) was stirred on an ice-bath. Mixture (1) was added dropwise. The mixture was stirred and refluxed overnight, then cooled and washed with water. The organic layer was separated, dried, filtered and the
35 solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 5.6 g of *N*-[4-[1-(phenylmethyl)-4-

piperidiny]phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 7, melting point 134°C).

EXAMPLE A.8

5 A mixture of intermediate (7) (0.025 mole) in methanol (250 ml) was hydrogenated at 50°C overnight with Pd/C 10% (2 g) as a catalyst. After uptake of hydrogen (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried. A part (0.2 g) of this fraction was purified by high performance liquid
10 chromatography over RP-18 (eluent: (NH₄OAc 0.5%/CH₃CN 90/10)/CH₃OH/CH₃CN 75/25/0, 0/50/50, 0/0/100 and 75/25/0; column: Hyperprep RP 100 Å 8 µm). The pure fractions were collected and the solvent was evaporated, yielding 0.119 g of *N*-[4-(4-piperidiny]phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide acetate (1:2) (intermediate 8,
15 melting point 150°C).

EXAMPLE A.9

A mixture of 4-[4-(phenylmethyl)-1-piperaziny]benzenamine (0.12 mole) in THF (300 ml) and triethylamine (50 ml) was stirred. [1,1'-Biphenyl]-2-carbonyl
20 chloride (0.12 mole) was added dropwise. The mixture was stirred overnight. The solvent was evaporated. The residue was dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE/2-propanol. The precipitate was filtered off and dried. A part (1 g) of this fraction was purified by column chromatography
25 over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1). The pure fractions were collected and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 0.84 g of *N*-[4-[4-(phenylmethyl)-1-piperaziny]phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 9, melting point 162°C).

30

EXAMPLE A.10

A mixture of intermediate (9) (0.1 mole) in methanol (500 ml) was hydrogenated for two hours with palladium-on-carbon (10%) (10 g) as a catalyst. After uptake of hydrogen (1 equivalent), the catalyst was filtered off and the filtrate was
35 evaporated. The residue was triturated in 2-propanol. The precipitate was filtered off and dried, yielding 29 g of *N*-[4-(1-piperaziny]phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 10, melting point 176°C).

EXAMPLE A.11

- a) A mixture of 4-(4-bromophenyl)-1-(phenylmethyl)-4-piperidinol (0.23 mole) and Cu₂O (2 g) in aqueous ammonia (500 ml) was stirred at 180°C for twelve hours. The mixture was cooled, extracted with DCM and washed with water. The organic layer was dried, filtered off and evaporated, yielding 60 g of 4-[1,2,3,6-tetrahydro-1-(phenylmethyl)-4-pyridinyl]benzenamine.
- b) [1,1'-Biphenyl]-2-carbonyl chloride (0.05 mole) was added dropwise to a stirring mixture of 4-[1,2,3,6-tetrahydro-1-(phenylmethyl)-4-pyridinyl]benzenamine (0.045 mole) in THF (300 ml) and triethylamine (25 ml). The mixture was stirred overnight. The solvent was evaporated. The residue was dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 18.5 g of N-[4-[1,2,3,6-tetrahydro-1-(phenylmethyl)-4-pyridinyl]phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 11, melting point 142°C).

EXAMPLE A.12

- A mixture of α -phenyl-4-piperidineacetonitrile hydrochloride (0.05 mole), 1-fluoro-4-nitrobenzene (0.06 mole) and potassium carbonate (0.15 mole) in DMF (200 ml) was stirred at 50°C for four hours. Water and DIPE were added. The mixture was cooled. The precipitate was filtered off, washed with water and DIPE and dried, yielding 11.6 g of (\pm)-1-(4-nitrophenyl)- α -phenyl-4-piperidineacetonitrile (intermediate 12, melting point 118°C).

EXAMPLE A.13

- A mixture of intermediate (12) (0.036 mole) in a 48% aqueous solution of hydrogen bromide (100 ml) was stirred and refluxed for three hours, cooled, poured out into water and extracted twice with DCM. The organic layer was separated, washed with water, dried, filtered and the solvent was evaporated. The residue was triturated with 2-propanol. The precipitate was filtered off and dried, yielding 9.5 g of (\pm)-1-(4-nitrophenyl)- α -phenyl-4-piperidineacetic acid (intermediate 13, melting point 216°C).

EXAMPLE A.14

- Thionyl chloride (0.01 mole) was added to a mixture of intermediate (13) (0.0029 mole) in DCM (10 ml). The mixture was stirred for one minute overnight and then allowed to stand. The solvent was evaporated. The residue was dissolved in DCM (10 ml). Methanol (10 ml) was added. The mixture was allowed to stand for four hours, then poured out into a NaHCO₃ solution and

extracted with DCM. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was triturated in hexane/DIPE. The precipitate was filtered off and dried, yielding 0.9 g of (±)-methyl 1-(4-nitrophenyl)-α-phenyl-4-piperidineacetate (intermediate 14, melting point 124°C).


EXAMPLE A.15


A mixture of intermediate (14) (0.0022 mole) in methanol (100 ml) was hydrogenated at 50°C with palladium-on-carbon (10%) (0.1 g) as a catalyst in the presence of a 4% thiophene solution (0.1 ml). After uptake of hydrogen (3 equivalents), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in hexane. The precipitate was filtered off and dried, yielding 0.7 g of (±)-methyl 1-(4-aminophenyl)-α-phenyl-4-piperidineacetate (intermediate 15, melting point 125°C).

EXAMPLES A.16 to A.18

In order to facilitate the understanding of these examples, the following presents a scheme (scheme 1) of the preparation of intermediate resins starting from a commercially available resin:

Scheme 1 :





R^c being ortho-iodophenyl, and Z^a being hydroxy or chloro.

EXAMPLE A.16

- 5 Ethylamine (0.0056 mole; 2.8 ml of a 2 M solution in THF, thus $R^a = a$ in scheme 1) was added to a Novabiochem 01-64-0261 commercial resin (1 g) in DCM (15 ml). Titanium (IV) isopropoxide (1.65 ml) was added and the mixture was shaken for one hour at room temperature. Triacetoxy borohydride (1.187 g) was added and the reaction mixture was shaken for 48 hours at room
- 10 temperature. The mixture was filtered, and the filter residue was washed three times with DCM, three times with a (1:1) mixture of DCM and methanol, three times with firstly methanol, then secondly with DCM, once with a mixture of 10 ml DCM and 2 ml DIPEA, three times with firstly DCM, then secondly with methanol, then dried, quantitatively yielding an ethylamino terminated resin
- 15 identified as I-a in scheme 1, which is used in the next reaction step without further purification.

EXAMPLE A.17

- Intermediate (13) (0.0056 mole) was added to the resin of example A.16
- 20 (0.00112 mole). A solution of a complex of (T-4)-hexafluorophosphate(1^-) (1-hydroxy-1H-benzotriazolato-O)tri-1-pyrrolidinyI-phosphorus(1^+) (hereinafter referred to as "PyBOP") (2.9 g) in DCM (15 ml) and DMF (5 ml) was added. Triethylamine (0.0112 mole) was added and the reaction mixture was shaken for 24 hours at room temperature, filtered and the filter residue was washed
- 25 with DMF (5 x 20 ml), then five times with (1:1) mixture of DCM and methanol (20 ml), five times with a (95:5) mixture of DCM with acetic acid (20 ml), five

times with DMF (20 ml) and three times with NMP (20 ml), quantitatively yielding a nitro terminated resin identified as II-a in scheme 1.

EXAMPLE A.18

- 5 A mixture of the resin of example A.17 (0.00112 mole) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (0.0224 mole) in NMP (20 ml) was shaken for six days at 55°C, then cooled, filtered and the filter residue was washed with DMF (three times), with a mixture of 10 ml DMF and 2 ml DIPEA, and then three times with firstly DCM, followed by secondly methanol, then dried, quantitatively yielding an amino terminated
10 resin identified as III-a in scheme 1.

EXAMPLE A.19

- Triisopropylamine (0.011 mole) was added to the resin of example A.18 (0.00112 mole) in DCM (10 ml). N,N-dimethyl-4-pyridinamine (0.0003 mole) in
15 DCM (3 ml) was added. A solution of o-iodobenzoyl chloride (0.00336 mole) in DCM (5 ml) was added and the reaction mixture was shaken overnight at room temperature. The mixture was filtered, the residue was washed three times with DCM, once with a mixture of DCM (10 ml) and DIPEA (2 ml), three times with DCM then methanol, then dried. The resulting product was treated with
20 benzylamine (1 ml) in DCM (10 ml) and shaken for 60 hours at room temperature. The mixture was filtered, washed three times with DCM, once with DCM/methanol 50/50, three times with DCM then methanol, then dried, yielding 0.00515 g (46%) of a resin identified as IV-a in scheme 1.

EXAMPLE A.20

The procedure of example A.16 is repeated while replacing ethylamine by n-propylamine, thus quantitatively yielding a n-propylamino terminated resin identified as I-b in scheme 1.

EXAMPLE A.21

The first experimental procedure of example A.17 is repeated while replacing the resin of example A.16 by the resin of example A.20, thus quantitatively yielding the resin identified as II-b in scheme 1.

EXAMPLE A.22

The procedure of example A.18 is repeated while replacing the resin of example A.17 by the resin of example A.21, thus quantitatively yielding the resin identified as III-b in scheme 1.

EXAMPLE A.23

The procedure of example A.19 is repeated while replacing the resin of example A.18 by the resin of example A.22, thus yielding the resin identified as IV-b in scheme 1.

5

EXAMPLE A.24

The procedure of example A.16 is repeated while replacing ethylamine by isopropylamine, thus quantitatively yielding an isopropylamino terminated resin identified as I-c in scheme 1.

10

EXAMPLE A.25

The first experimental procedure of example A.17 is repeated while replacing the resin of example A.16 by the resin of example A.24, thus quantitatively yielding the resin identified as II-c in scheme 1.

15

EXAMPLE A.26

The procedure of example A.18 is repeated while replacing the resin of example A.17 by the resin of example A.25, thus quantitatively yielding the resin identified as III-c in scheme 1.

20

EXAMPLE A.27

The procedure of example A.19 is repeated while replacing the resin of example A.18 by the resin of example A.26, thus yielding the resin identified as IV-c in scheme 1.

25

EXAMPLE A.28

The procedure of example A.16 is repeated while replacing ethylamine by phenylamine, thus quantitatively yielding a phenylamino terminated resin identified as I-d in scheme 1.

30

EXAMPLE A.29

The first experimental procedure of example A.17 is repeated while replacing the resin of example A.16 by the resin of example A.28, thus quantitatively yielding the resin identified as II-d in scheme 1.

35

EXAMPLE A.30

The procedure of example A.18 is repeated while replacing the resin of example A.17 by the resin of example A.29, thus quantitatively yielding the resin identified as III-d in scheme 1.

40

EXAMPLE A.31

The procedure of example A.19 is repeated while replacing the resin of example A.18 by the resin of example A.30, thus yielding the resin identified as IV-d in scheme 1.

EXAMPLE A.32

The procedure of example A.16 is repeated while replacing ethylamine by benzylamine, thus quantitatively yielding a benzylamino terminated resin identified as I-e in scheme 1.

EXAMPLE A.33

The first experimental procedure of example A.17 is repeated while replacing the resin of example A.16 by the resin of example A.32, thus quantitatively yielding the resin identified as II-e in scheme 1.

EXAMPLE A.34

The procedure of example A.18 is repeated while replacing the resin of example A.17 by the resin of example A.33, thus quantitatively yielding the resin identified as III-e in scheme 1.


EXAMPLE A.35


The procedure of example A.19 is repeated while replacing the resin of example A.18 by the resin of example A.34, thus yielding the resin identified as IV-e in scheme 1.

EXAMPLES A.36 to A.38

In order to facilitate the understanding of these examples, the following presents another scheme (scheme 2) for the preparation of intermediate resins starting from a commercially available resin :

Scheme 2 :



**EXAMPLE A.36**

A mixture of commercial Novabiochem 01-64-0261 resin (25.1 g, 0.028 mole),
 5 4-bromoaniline (24 g, 0.140 mole) and titanium (IV) isopropoxide (41 ml, 0.140 mole) in DCM (400 ml) was stirred gently for one hour at room temperature. Sodium triacetoxyborohydride (30 g, 0.140 mole) was added and the reaction mixture was stirred overnight at room temperature. Methanol (50 ml) was added and the mixture was stirred for one hour, then filtered, washed once with
 10 DCM, once with methanol, then once with DCM (200 ml) + DIPEA (20 ml), washed three times with firstly DCM, followed secondly by methanol, then dried, yielding 29.28 g of a resin identified as V in scheme 2, which is used in the next reaction step without further purification.

EXAMPLE A.37

4-Phenyl benzoic acid (8.3 g, 0.042 mole) was dissolved in DCM (100 ml). Thionyl chloride (10 g, 0.084 mole) was added. DMF (10 drops) was added and the mixture was stirred and refluxed for one hour. The solvent was evaporated. DCM (three times 50 ml) was added. The solvent was evaporated.
 20 The residue was dissolved in DCM (50 ml). This solution was added to a mixture of the resin of example A.36 (14.64 g, 0.0133 mole), DIPEA (24 ml, 0.140 mole) and 4-dimethylaminopyridine (hereinafter referred as DMAP) (0.5 g) in DCM (150 ml). The reaction mixture was shaken overnight at room temperature, then filtered and the filter residue was washed with 100 ml DMF +
 25 20 ml DIPEA, then with methanol, water, DCM, methanol, DCM and methanol, and dried, yielding 15.73 g of a resin identified as VI-a in scheme 2.

EXAMPLE A.38

- 4'-(Trifluoromethyl)-2-biphenyl carboxylic acid (14.64 g, 0.042 mole) was dissolved in DCM (100 ml). DMF (1 ml) was added. Thionyl chloride (10 g, 0.084 mole) was added and the mixture was stirred and refluxed for one hour.
- 5 The solvent was evaporated. DCM (twice 50 ml) was added, then the solvent was evaporated. The residue was dissolved in DCM (50 ml). This solution was added to a mixture of the resin of example A.36 (14.64 g, 0.0133 mole), DIPEA (24 ml, 0.140 mole) and DMAP (0.5 g) in DCM (150 ml). The reaction mixture was shaken for four hours at room temperature then filtered and the filter
- 10 residue was washed with 100 ml DMF + 20 ml DIPEA, then washed three times firstly with DCM and secondly with methanol, and finally dried. This reaction product was reacted once more with half the initial quantities of 4'-(trifluoromethyl)-2-biphenyl carboxylic acid, thionyl chloride, DIPEA and DMAP. The reaction mixture was shaken overnight at room temperature, then filtered, and
- 15 the filter residue was shaken with DMF + 20 ml DIPEA, then methanol, water, methanol, DCM, methanol, DCM and methanol, then dried, yielding 17.48 g of a resin identified as VI-b in scheme 2.

Example A.39

- 20 a) A mixture of 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.09 mol) in DCM (500 ml) and DMF (5 ml) was stirred. Ethanedioyl dichloride (0.09 mol) was added dropwise. The mixture was stirred for 1 hour to give mixture 1. A mixture of 4-[1-(phenylmethyl)-4-piperidinyl]-benzenamine hydrochloride salt (1:1) (0.046 mol) in DCM (500 ml) and triethylamine (20 ml) was stirred on an
- 25 ice-bath. Mixture 1 was added dropwise. The mixture was stirred and refluxed overnight, then cooled and washed with water. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected and the solvent was evaporated. The
- 30 residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 5.6 g of N-[4-[1-(phenylmethyl)-4-piperidinyl]phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 16, mp. 134°C).
- b) A mixture of intermediate (16) (0.025 mol) in methanol (250 ml) was hydrogenated at 50°C overnight with Pd/C 10% (2 g) as a catalyst. After uptake
- 35 of hydrogen (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 7.7g N-[4-(4-piperidinyl)phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 17).

Example A.40

- a) [1,1'-Biphenyl]-2-carboxylic acid (0.25 mol) was dissolved in DCM (500 ml) and DMF (0.5 ml). Thionyl chloride (0.51 mol) was added dropwise. The mixture was stirred and refluxed for 1 hour under nitrogen flow. The solvent was evaporated. DCM (500 ml) was added twice. The solvent was evaporated twice. The residue was dissolved in DCM (200 ml) and then added dropwise at 0°C to a mixture of 4-[1-(phenylmethyl)-4-piperidiny]-benzenamine (0.25 mol) and *N*-(1-methylethyl)-2-propanamine (0.75 mol) in DCM (800 ml). The mixture was brought to room temperature and then stirred at room temperature overnight under nitrogen flow. The mixture was washed three times with water (800 ml). The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 125 g of *N*-[4-[1-(phenylmethyl)-4-piperidiny]phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 18).
- b) A mixture of intermediate (18) (0.145 mol) in methanol (500 ml) was hydrogenated at 50°C during 48 hours with Pd/C (10%, 3 g) as a catalyst. After uptake of hydrogen (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 49 g of *N*-[4-(4-piperidiny)phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 19).

Example A.41

- a) 4'-(Trifluoromethyl)-[1,1'-biphenyl]-2-carbonyl chloride (0.12 mol) was added dropwise to a stirring mixture of 4-[1,2,3,6-tetrahydro-1-(phenylmethyl)-4-pyridiny]benzenamine (0.095 mol) in DCM (300 ml) and triethylamine (50 ml). The mixture was stirred overnight, poured out into water and then stirred for 30 minutes. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 43 g of *N*-[4-[1,2,3,6-tetrahydro-1-(phenylmethyl)-4-pyridiny]phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 20).
- b) 1-Chloroethyl chloroformate (0.078 mol) was added dropwise to a stirring mixture of intermediate (20) (0.039 mol) in 1,2-dichloro-ethane (500 ml). The mixture was stirred for 30 minutes and then stirred and refluxed overnight. The solvent was evaporated. Methanol (500 ml) was added. The mixture was stirred and refluxed overnight. The solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 20.8 g of *N*-[4-(1,2,3,6-tetrahydro-4-pyridiny) phenyl]-4'-(trifluoro-methyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 21).

Example A.42

A mixture of intermediate (11) (0.04 mol) in 1,2-dichloro-ethane (200 ml) was stirred on an ice-bath. 1-Chloroethyl chloroformate (15 ml) was added dropwise at a temperature below 5°C. The mixture was stirred for 1 hour and then stirred
5 and refluxed overnight. The solvent was evaporated. Methanol (200 ml) was added. The mixture was stirred and refluxed for 2 hours. The solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yield 16.7g of *N*-[4-(1,2,3,6-tetrahydro-4-pyridinyl)phenyl]-[1,1'-biphenyl]-2-carboxamide (intermediate 22).

10

Example A.43

a) A mixture of intermediate (6) (0.0047 mol) and di-tert-butyl dicarbonate (0.0052 mol) in DCM (50ml) was stirred at room temperature for 2 hours. DMF (5 ml) was added. The mixture was stirred at room temperature for 3 hours.
15 The solvent was evaporated. The residue was stirred in DIPE. The precipitate was filtered off and dried in vacuo at 54°C, yielding 2.47g of 1,1-dimethylethyl ester 4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazinecarboxylic acid (intermediate 23, mp. 204°C).


b) NaH 60% in mineral oil (0.0056 mol) was treated with hexane; stirred under
20 nitrogen flow and decanted. DMF dry (25ml) was added to the residue. The suspension was stirred at room temperature under nitrogen flow. A solution of intermediate (23) (0.00375 mol) in DMF dry (25 ml) was added dropwise. The mixture was stirred at room temperature under nitrogen flow for 2.5 hours. A solution of methyl methanesulfonate (0.0045 mol) in DMF dry (50 ml) was
25 added dropwise. The mixture was stirred at room temperature for 18 hours. Water (150 ml) was added. The mixture was extracted twice with DCM. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was stirred in 60 ml of hexane/DIPE (3:1). The mixture was stirred and refluxed until a clear solution was obtained and then brought to room
30 temperature. The precipitate was filtered off and dried, yielding 1.6 g of 1,1-dimethylethyl ester 4-[4-[methyl[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazinecarboxylic acid (intermediate 24).

c) A solution of trifluoroacetate (20 ml) and DCM (200 ml) was added to intermediate (24) (0.0024 mol) and stirred for 1 hour and 30 minutes at room
35 temperature. The solvent was evaporated. DCM was added and again the solvent was evaporated, yielding 1.7 g of *N*-methyl-*N*-[4-(1-piperazinyl)phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (intermediate 25).

Example A.44

Intermediate resins wherein X_2 and X_3 represent N and Z_1 and Z_2 represent $-\text{CH}_2\text{CH}_2-$ can be prepared as depicted in scheme 3 starting from a commercially available resin :


5

Scheme 3 :


10

a) A mixture of commercial Novabiochem 01-64-0261 resin (0.180 g, 0.002 mol), 4-(1-tert-butoxycarbonylpiperazin-4-yl)aniline ((0.001 mol), dissolved in

- DCM (2 ml)) and titanium (IV) isopropoxide (0.001 mol) in DCM (3 ml) was shaken for 2 hours at room temperature. Sodium triacetoxyborohydride (0.001 mol) was added portionwise and the reaction mixture was shaken for 20 hours at room temperature. The mixture was filtered and the filter residue was washed three times with DCM (3 times), methanol (3 times), and then three times with DCM, then dried, yielding resin (VII-a).




- b) 2-Bromo-4-methylbenzoic acid (0.001 mol) in DCM (5 ml) with thionyl chloride (0.013 mol) was stirred and refluxed for one hour. The mixture was blown dry under nitrogen. Again, DCM (5 ml) with thionyl chloride (0.013 mol) was added. The reaction mixture was stirred and refluxed for one hour. The mixture was blown dry under nitrogen and three times with DCM (3 ml) was added, then evaporated again. The residue was dissolved in DCM (3 ml), this solution was added to a solution of resin (VII-a) (0.0002 mol) in DCM (1 ml). DMAP (0.0002 mol) in DCM (1 ml) was added. DIPEA (0.002 mol) was added and the reaction mixture was shaken for 20 hours at room temperature. The reaction mixture was filtered and the filter residue was washed three times with DCM (3 times), methanol (3 times), and then three times with DCM, then dried, yielding resin (VIII-a).



- c) A solution of 4-(trifluoromethyl)benzeneboronic acid (0.0016 mol) in dioxane (3 ml) was added to resin (VIII-a) (0.0002 mol) which was previously washed with dioxane (5 ml). Then, KOH (0.0032 mol of a 2 M solution) was added and the reaction mixture was shaken for 30 minutes at room temperature under a nitrogen atmosphere. $\text{PdCl}_2(\text{PPh}_3)_2$ (0.00004 mol) in NMP (0.5 ml) was added and the reaction mixture was shaken for 2 hours at 90°C. Again, $\text{PdCl}_2(\text{PPh}_3)_2$

(0.00004 mol) in NMP (0.5 ml) was added and the reaction mixture was shaken for 2 hours at 90°C. The mixture was cooled, filtered and the filter residue was washed with DMF (3 times), with water (3 times), DMF (3 times), methanol (3 times), DCM (3 times), methanol (3 times) and DCM (3 times), then dried yielding a residue. Said residue was stirred in a solution of TMSTf (1 M) and 2,6-lutidine (1.5 M) in DCM (4 ml) for 2 hours at room temperature, filtered, and washed with DCM (3 times) and methanol (3 times), and dried, yielding resin (IX-a).




10

Example A.45


a) A suspension of 4-(*tert*-butoxycarbonylamino)piperidine (15 equivalents) in NMP (2 ml) was added to resin (VI-b) in NMP (1 ml). [1,1'-Binaphthalene]-2,2'-diylbis[diphenyl-phosphine (BINAP) (0.00011 mol) was added portionwise. *tert*-Butoxysodium (15 equivalents) was added portionwise. The reaction mixture was shaken for one hour under nitrogen flow. Pd₂(dba)₃ (0.000022 mol) in NMP (1 ml) was added and the reaction mixture was shaken for 18 hours at 105°C. The mixture was cooled, filtered and the filter residue was washed with three times (with DMF followed with water), and then three times with methanol, then DCM, yielding resin (X).

20



- b) Resin (X) (0.00011 mol) was shaken in NMP (2 ml). Bromobenzene (0.00165 mol) in NMP (1 ml) was added. BINAP (0.068 g) was added portionwise. *Tert*-butoxysodium (0.190 g) was added portionwise. The mixture was shaken for 1 hour under nitrogen. $\text{Pd}_2(\text{dba})_3$ (0.020 g) in NMP (1 ml) was added and the reaction mixture was shaken for 18 hours at 105°C, then cooled, filtered and the filter residue was washed with three times (with DMF followed with water), and then three times with methanol, then DCM yielding resin (XI).


10



Example A.45

- Resin (XI) (0.0002 mol) was washed with DCM (4 ml), then filtered off, then dissolved in DCM (5 ml). Trichloromethyl chloroformate (0.001 mol) was added and the reaction mixture was shaken for 4 hours at room temperature. The mixture was filtered, washed three times with DCM, then dried, yielding resin (XII).

15



Example A.46

- 282203 a) 1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (EDCI)
 5 (0.096 mol) was added at room temperature to a mixture of 4-(4-aminophenyl)-
 1-piperazinecarboxylic acid ethyl ester (0.08 mol), 2-iodo benzoic acid
 (0.096 mol) and 1-hydroxybenzotriazole (HOBT) (0.096 mol) in DCM (500 ml).
 The mixture was stirred at room temperature overnight. Water was added. The
 mixture was extracted with DCM. The organic layer was separated, dried,
 10 filtered, and the solvent was evaporated. The residue was taken up in DIPE.
 The precipitate was filtered off and dried, yielding 39 g of 4-[4-[(2-iodobenzoyl)-
 amino]phenyl]-1-piperazinecarboxylic acid, ethyl ester (intermediate 27).
 b) A mixture of intermediate (27) (0.041 mol) and potassium hydroxide
 (0.41 mol) in isopropanol (200 ml) was stirred and refluxed for 3 hours and the
 15 solvent was evaporated till dryness. Water was added. The mixture was
 extracted with DCM and the solvent was evaporated, yielding 2-iodo-N-[4-(1-
 piperaziny]phenyl]benzamide (intermediate 28).
 c) α -Bromo-benzeneacetic acid methyl ester (0.0123 mol) was added at room
 temperature to a mixture of intermediate (28) (0.0123 mol) and Na_2CO_3
 20 (0.0123 mol) in DMF (50 ml). The mixture was stirred at room temperature for
 2 hours. Water was added. The mixture was stirred at room temperature for
 15 minutes. The precipitate was filtered, washed with diethyl ether and dried,
 yielding 5.8 g of 4-[4-[(2-iodobenzoyl)amino]phenyl]- α -phenyl-1-piperazine-
 acetic acid, methyl ester (intermediate 29).
 25 d) A mixture of intermediate (29) (0.0036 mol), tributyl-2-furanyl-stannane
 (0.029 mol), $\text{PdCl}_2(\text{PPh}_3)_2$ (0.0007 mol) and Na_2CO_3 (0.0576 mol) in dioxane
 (30 ml) was stirred and refluxed for 1 hour. Water was added. The mixture
 was extracted with ethyl acetate. The organic layer was separated, dried,
 filtered, and the solvent was evaporated. The residue was purified by column
 30 chromatography over silica gel (eluent : cyclohexane/ethyl acetate 80/20). The

pure fractions were collected and the solvent was evaporated, yielding 4-[4-[[2-(2-furanyl)benzoyl]amino]phenyl]- α -phenyl-1-piperazineacetic acid, methyl ester (intermediate 30, mp. 90°C).

- 5 e) A mixture of intermediate (30) (0.0006 mol) and potassium hydroxide (0.006 mol) in isopropanol (5 ml) was stirred at room temperature overnight. The solvent was evaporated. The residue was dissolved in isopropanol/HCl 6N and converted into the hydrochloric acid salt. The precipitate was filtered off and dried, yielding 0.31 g of 4-[4-[[2-(2-furanyl)benzoyl]amino]phenyl]- α -phenyl-1-piperazineacetic acid (intermediate 31).

10

B. preparation of polyarylcaboxamide compounds of the invention.

EXAMPLE B.1

- A mixture of 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.012 mole) in DCM (100 ml) and DMF (8 drops) was stirred. Ethanedioyl dichloride (0.012
15 mole) was added. The mixture was stirred for 2 hours, to give mixture (I). Triethylamine (8 ml) was added to a mixture of intermediate (4) (0.005 mole) in DCM (100 ml). The mixture was stirred on an ice-salt bath to give mixture (II). Mixture (I) was added dropwise to mixture (II) and the resulting reaction mixture was stirred and refluxed for two days. The solvent was evaporated. The residue
20 was dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 99/1). The desired fractions were collected and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 2.99 g of
25 *N*-[4-[4-[2-oxo-1-phenyl-2-[(2,2,2-trifluoroethyl)amino]ethyl]-1-piperazinyl]-phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (melting point 208°C) identified as compound No. 1 in the following table F-1.

EXAMPLE B.2

- 30 Fluorene-4-carboxylic acid (0.00032 mole) in 1 ml of 1/1-mixture of DCM/NMP was added to PyBOP (0.00064 mole) in DCM (1 ml). This mixture was stood for 30 minutes, then added to the resin of example A.34. DCM (5 ml) was added, followed by DIPEA (0.00085 mole). The reaction mixture was shaken for 24 hours at room temperature, then filtered, washed 3 times with DCM, 3 times
35 with methanol, followed by DCM. A mixture of TFA/DCM/TIS (5/93/2) (4 ml) was added and the mixture was shaken for one hour at room temperature. The mixture was filtered, the filter residue was washed with a mixture of TFA/DCM/TIS (5/93/2) (2 ml) and with DCM (2 ml). The filtrate was blown dry at 50°C under a gentle stream of nitrogen gas, dissolved in DCM (5 ml) and DMF

(1 ml) and a Novabiochem 01-64-0171 resin was added while stirring the mixture at room temperature. Then, after one hour, an Argonaut P/N 800277 resin (0.040 g) was added. The mixture was stirred for four hours at room temperature, filtered, and the filtrate was blown dry at 50°C under a nitrogen flow, yielding 0.027 g of the compound identified as No. 2 in the following table F-1.

Compounds identified as No. 3 to No.11 in the following table F-1 were similarly prepared while using the same experimental procedure and replacing fluorene-4-carboxylic acid by the appropriate reactive acid.

EXAMPLE B.3

Fluorene-4-carboxylic acid (0.00028 mole) in 1 ml of a 1/1 mixture of DCM/NMP was added to PyBOP (0.00028 mole) in DCM (1 ml). This mixture was stood for 30 minutes, then added to the resin of example A.22. DCM (5 ml) was added, followed by triethylamine (0.00057 mole). The reaction mixture was shaken for 20 hours at room temperature, then filtered, washed three times with DCM, three times with first methanol, followed by secondly DCM. 4 ml of a mixture of TFA/DCM/TIS (5/93/2) was added and the mixture was shaken for one hour at room temperature. The mixture was filtered, the filter residue was washed with a mixture of TFA/DCM/TIS (2 ml; 5/93/2) and with DCM (1 ml). The filtrate was blown dry at 50°C under a gentle stream of nitrogen. This fraction was purified by high performance liquid chromatography over Hyperprep RP-C18 BDS (100 g, 100 Å, 8 µm; eluent: [(0.5% NH₄OAc in H₂O)/CH₃CN 90/10])/CH₃OH/CH₃CN (0 minutes) 75/25/0, (10 minutes) 0/50/50, (16 minutes) 0/0/100, (18.10-20.00 minutes) 75/25/0). The pure fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with DCM/aqueous K₂CO₃ solution, then separated over Extrelut™. The organic phase was blown dry under nitrogen at 50°C. The residue was dried further under vacuum at 60°C, yielding 0.002 g of the compound identified as No. 12 in the following table F-1.

Compounds identified as No. 13 to No. 20 in the following table F-1 were similarly prepared while using the same experimental procedure and replacing fluorene-4-carboxylic acid by the appropriate reactive acid.

EXAMPLE B.4

Fluorene-4-carboxylic acid (0.00015 mole) in 1 ml of 1/1 mixture of DCM/NMP was added to PyBOP (0.0003 mole) in DCM (1 ml). This mixture was stood for 30 minutes, then added to the resin of example A.26. DCM (5 ml) was added,

followed by DIPEA (0.00057 mole). The reaction mixture was shaken for 24 hours at room temperature, then filtered, washed three times with DCM, three times with first methanol, followed by secondly DCM. 4 ml of a mixture of TFA/DCM/TIS (5/93/2) was added and the mixture was shaken for one hour at room temperature. The mixture was filtered, the filter residue was washed with 2 ml of a mixture of TFA/DCM/TIS (5/93/2) and with DCM (1 ml). The filtrate was blown dry at 50°C under a gentle stream of nitrogen. This fraction was purified by high performance liquid chromatography over Hyperprep C18 BDS (100 g, 100 Å, 8 µm; eluent: [(0.5% NH₄OAc in H₂O)/CH₃CN 90/10)]/CH₃OH/CH₃CN (0 minute) 75/25/0, (10 minute) 0/50/50, (16 minute) 0/0/100, (18.10-20.00 minute) 75/25/0). The pure fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with DCM/aqueous potassium carbonate solution, then separated over ExtrelutTM. The organic phase was blown dry under nitrogen at 50°C. The residue was dried further under vacuum at 60 °C, yielding 0.002 g of the compound identified as No. 21 in the following table F-1.

Compounds identified as No. 22 to No. 28 in the following table F-1 were similarly prepared while using the same experimental procedure and replacing fluorene-4-carboxylic acid by the appropriate reactive acid.

EXAMPLE B.5

Fluorene-4-carboxylic acid (0.00023 mole) in 1 ml of 1/1 mixture of DCM/NMP was added to PyBOP (0.00046 mole) in DCM (1 ml). This mixture was stood for 30 minutes, then added to the resin of example A.30. DCM (5 ml) was added, followed by DIPEA (0.00057 mole). The reaction mixture was shaken for 24 hours at room temperature, then filtered, washed three times with DCM, three times with first methanol, followed by secondly DCM. A mixture of TFA/DCM/TIS (4 ml; 75/23/2) was added and the mixture was shaken for one hour at room temperature. The mixture was filtered, the filter residue was washed with 2 ml of a mixture of TFA/DCM/TIS (75/23/2) and with DCM (2 ml). The filtrate was blown dry at 50°C under a gentle stream of nitrogen. The residue was dissolved in DCM (5 ml), then blown dry once more. This fraction was purified by high performance liquid chromatography over Hyperprep RP-C18 BDS (100 g, 100 Å, 8 µm; eluent: [(0.5% NH₄OAc in H₂O)/CH₃CN 90/10)]/CH₃OH/CH₃CN (0 minute) 75/25/0, (10 minutes) 0/50/50, (16 minutes) 0/0/100, (18.10-20.00 minutes) 75/25/0). The pure fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with DCM/aqueous potassium carbonate solution, then separated over

Extrelut™. The organic phase was blown dry under nitrogen at 50°C. The residue was dried further under vacuum at 60°C, yielding 0.0046 g of the compound identified as No. 29 in the following table F-1.

Compounds identified as No. 30 to No. 36 in the following table F-1 were
5 similarly prepared while using the same experimental procedure and replacing fluorene-4-carboxylic acid by the appropriate reactive acid.

EXAMPLE B.6

A mixture of the Novabiochem 01-64-0261 commercial resin (0.00011 mole),
10 intermediate (15) (0.00061 mole) and isopropyl titanate (0.18 ml) in DCM (5 ml) was shaken for one hour at room temperature. Triacetoxy borohydride (0.128 g) was added and the reaction mixture was shaken for 16 hours at room temperature. Methanol (1 ml) was added and the mixture was shaken for 5 minutes, then filtered, washed three times with DCM, methanol and dried under
15 vacuum, yielding a residue (1).

Thionyl chloride (0.0020 mole) was added to 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.00055 mole) in DCM (2 ml) and the mixture was refluxed for 30 minutes while stirring followed by evaporation of the solvent. The residue was dissolved in DCM (5 ml) and the solution was added to the above-prepared
20 residue (1). DIPEA (0.0011 mole) was added, followed by the addition of N,N-dimethyl-4-pyridinamine (0.00008 mole). The reaction mixture was shaken for 21 hours at room temperature, then filtered, washed three times with DCM, then three times with first a 4% acetic acid/DCM mixture, secondly DCM and finally three times with first DCM and secondly methanol, then dried, and 4 ml
25 of a mixture of TFA/DCM/TIS (49/49/2) was added. This mixture was shaken for one hour, filtered, washed with 2 ml of a mixture of TFA/DCM/TIS (49/49/2) and DCM (2 ml). The filtrate was blown dry with nitrogen at 50°C. DCM (5 ml) was added, then removed under a nitrogen stream at 50°C, yielding 0.080 g of methyl α -phenyl-1-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]
30 carbonyl]amino]phenyl]-4-piperidineacetate trifluoroacetate (1:1) identified as compound No. 37 in the following table F-1.

EXAMPLE B.7

4'-(Trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.00033 mole) and PyBOP
35 (0.171 g) were dissolved in DCM (5 ml). This mixture was added to the resin of example A.22 (0.00066 mole). DIPEA (0.00066 mole) was added and the reaction mixture was shaken for 48 hours at room temperature, filtered and the residue was washed three time with DMF, then three times with DCM and methanol, dried, yielding a residue. Said residue and a TFA/DCM/TIS (5:93:2)

- (4 ml) was shaken for 30 minutes at room temperature, then filtered, washed with a mixture of TFA/DCM/TIS (5/93/2) (2 ml) and DCM (2 ml), then the filtrates were blown dry with nitrogen gas at 50°C, then dried further under vacuum at 60°C, yielding 0.030 g of *N*-[4-[4-[2-oxo-1-phenyl-2-(propylamino)ethyl]-1-piperidinyl] phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide trifluoroacetate (1:1) identified as compound No. 38 in the following table F-1.

EXAMPLE B.8

- DMF (0.5 ml) was added to a solution of 4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxylic acid (0.014 mole) in DCM (50 ml) and thionyl chloride (0.028 mole). The mixture was stirred and refluxed for one hour. The solvent was evaporated. DCM (50 ml) was added twice and the solvent was evaporated. The residue was dissolved in DCM (20 ml) and this solution was added to a mixture of intermediate (15) (0.014 mole) and DIPEA (0.028 mole) in DCM (80 ml). The mixture was stirred at room temperature for three hours and washed with water. The organic layer was dried and the solvent was evaporated. The residue was crystallized from 2-propanol. The precipitate was filtered off and dried, yielding 6.2 g of methyl α -phenyl-1-[4-[[[4'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-4-piperidineacetate (melting point 151°C) identified as compound No. 39 in the following table F-1.

EXAMPLE B.9

- A mixture of intermediate (6) (0.023 mole) and Na₂CO₃ (0.023 mole) in DMF (150 ml) was stirred. Methyl α -bromo- α -phenylacetate (0.023 mole) was added dropwise. The mixture was stirred overnight. The solvent was evaporated. The residue was dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 11.4 g of methyl α -phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazineacetate identified as compound No. 40 in the following table F-1.

EXAMPLE B.10

- A mixture of intermediate (8) (0.018 mole) and Na₂CO₃ (0.03 mole) in DMF (100 ml) was stirred. Methyl α -bromo- α -phenylacetate (0.025 mole) was added dropwise. The mixture was stirred overnight. The solvent was evaporated. The residue was dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1). The pure

fractions were collected and the solvent was evaporated, yielding 7.2 g of methylphenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperidineacetate (melting point 138°C) identified as compound No. 41 in the following table F-1.

5

EXAMPLE B.11

Methyl α -bromo- α -phenylacetate (0.1 mole) was added dropwise to a stirring mixture of intermediate (10) (0.07 mole) and Na_2CO_3 (13 g) in DMF (300 ml). The mixture was stirred overnight. The solvent was evaporated. The residue was crystallized from methanol. The precipitate was filtered off and dried, yielding 30.2 g of methyl 4-[4-[[[1,1'-biphenyl]-2-ylcarbonyl]amino]phenyl]- α -phenyl-1-piperazineacetate (melting point 125°C) identified as compound No. 52 in the following table F-1.

15 EXAMPLE B.12

- a) A mixture of compound (40) (0.19 mole) in HCl (36%) (100 ml) was stirred and refluxed for five hours, then stirred overnight at room temperature. The precipitate was filtered off and triturated under 2-propanol, filtered off and dried, yielding 5 g of the intermediate compound α -phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl] piperazineacetic acid monohydrochloride.
- b) A mixture of the intermediate compound obtained in step (a) (0.00016 mole), PyBOP (0.00032 mole) and triethylamine (0.1 ml) in DCM (5 ml) was stirred for 30 minutes. Ethylamine (0.0005 mole) was added and the reaction mixture was stirred overnight at 40°C. The reaction mixture was cooled. Water (2 ml) was added and the mixture was stirred for 15 minutes, then filtered through ExtrelutTM, and the desired compound was isolated by high performance liquid chromatography, yielding 0.046 g of *N*-[4-[4-[2-(ethylamino)-2-oxo-1-phenylethyl]-1-piperazinyl]phenyl]-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide (melting point 123°C) identified as compound No. 54 in the following table F-1.

Compounds identified as No. 55 to No. 61 in the following table F-1 were similarly prepared while using the same experimental procedure and replacing ethylamine by the appropriate reactive amine.

35

EXAMPLE B.13

A mixture of α -phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazineacetic acid monohydrochloride (0.011 mole) in sulfuric acid (10 ml) and propanol (150 ml) was stirred and refluxed overnight.

The solvent was evaporated. The residue was dissolved in DCM and washed with a solution of Na_2CO_3 . The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{CN}$ 95/5). The pure
5 fractions were collected and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 2.6 g of propyl-phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazineacetate (melting point 151°C) identified as compound No. 62 in the following table F-1.

10

EXAMPLE B.14

A mixture of intermediate (6) (0.017 mole) and ethyl 2-phenylacrylate (0.017 mole) in DMF (100 ml) was stirred for two days. Na_2CO_3 (1 g) was added. The mixture was stirred for two days. The solvent was evaporated. The residue was
15 dissolved in DCM. The organic layer was separated, washed, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 10.6 g of ethylphenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]-phenyl]-1-piperazine propanoate (melting point 195°C) identified as compound No. 81 in the
20 following table F-1.

EXAMPLE B.15

a) A mixture of compound No. 81 (0.016 mole) in HCl (36%) (100 ml) was stirred and refluxed for eight hours, then cooled and filtered. The residue was
25 triturated in 2-propanol. The precipitate was filtered off and dried. A part (0.2 g) of this fraction was purified by high performance liquid chromatography over RP-18 eluent: (NH_4OAc 0.5%/ CH_3CN 90/10)/ $\text{CH}_3\text{OH}/\text{CH}_3\text{CN}$ 75/25/0, 0/50/50 and 75/25/0; column: HYPERPREP 8 μm). The pure fractions were collected and the solvent was evaporated. The residue was triturated in DIPE.
30 The precipitate was filtered off and dried, yielding 0.12g of the intermediate compound 2-phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperazinepropanoic acid (melting point 202°C).
b) A mixture of the intermediate compound obtained in step (a) (0.00016 mole), PyBOP (0.00032 mole) and triethylamine (0.1 ml) in DCM (5 ml) was stirred
35 for 30 minutes. Propylamine (0.0004 mole) was added and the reaction mixture was stirred overnight at 40°C . The reaction mixture was cooled, washed with water (2 ml), then filtered through ExtrelutTM, and the extract's solvent was evaporated. The desired compound was isolated by high performance liquid chromatography over Hyperprep RP-C18 BDS (100 g,

100 Å, 8 µm; eluent : [(0.5% NH₄OAc in H₂O)/CH₃CN 90/10)]/CH₃OH/CH₃CN
(0 min) 75/25/0, (10 min) 0/50/50, (16 min) 0/0/100, (18.10-20.00 min) 75/25/0).

The pure fractions were collected and the solvent was evaporated, yielding of
N-[4-[4-[3-oxo-2-phenyl-3-(propylamino)propyl]-1-piperazinyl]phenyl]-4'-

5 (trifluoromethyl)-[1,1'-biphenyl]-2-carboxamide identified as compound No. 63 in
the following table F-1.

Compounds identified as No. 64 to No. 67 in the following table F-1 were
similarly prepared while using the same experimental procedure and replacing
propylamine by the appropriate reactive amine.

10

EXAMPLE B.16

a) A mixture of compound No. 41 (0.012 mole) in HCl (36%) (100 ml) was
stirred and refluxed for six hours, then stirred overnight at room temperature.

15 The precipitate was filtered off and triturated under 2-propanol. The
precipitate was filtered off and dried, yielding 6.2 g of the intermediate
compound α-phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]
amino] phenyl]-1-piperidineacetic acid monohydrochloride.

b) A mixture of the intermediate compound obtained in step (a) (0.00017 mole),
PyBOP (0.3 g) and triethylamine (0.1 ml) in DCM (5 ml) was stirred for 30
20 minutes. Ethylamine (0.00017 mole) was added. The reaction mixture was
stirred overnight at 40°C, then cooled and water (2 ml) was added. The
mixture was stirred for one hour, then filtered through Extrelut™, and the
filtrate was evaporated. The residue was purified by column chromatography
over silica gel (eluent : CH₂Cl₂/CH₃OH 90/10). The product fractions were
25 collected and the solvent was evaporated, yielding 0.010 g of *N*-[4-[1-[2
-(ethylamino)-2-oxo-1-phenylethyl]-4-piperidinyl]phenyl]-4'-(trifluoromethyl)
-[1,1'-biphenyl]-2-carboxamide identified as compound No. 69 in the
following table F-1.

Compounds identified as No. 70 to No. 80 in the following table F-1 were
30 similarly prepared while using the same experimental procedure and replacing
ethylamine by the appropriate reactive amine.

EXAMPLE B.17

The resin of example A.23 (0.000045 mole) was washed twice with dioxane.
35 1,4-Dioxane (1 ml) was added. 2,4-Difluorophenylboronic acid (0.0004 mole) in
1,4-dioxane (1 ml) was added. KOH (2 M) (0.25 ml) was added. The mixture
was shaken for 30 minutes under an argon atmosphere. PdCl₂(PPh₃)₂
(0.00001 mole) in NMP (0.250 ml) was added. The mixture was stirred for 90
minutes at 98°C. Again, PdCl₂(PPh₃)₂ was added and the reaction mixture was

warmed for 90 minutes at 98°C. The mixture was allowed to cool to room temperature, then filtered and the filter residue was washed three times with dioxane, three times with water and three times with methanol, then three times with DCM then methanol, finally three times with DCM. F (4 ml) was added. The mixture was shaken for 30 minutes, filtered, washed with TFA/DCM/TIS (2 ml, 5/93/2) and DCM (2 ml), and the filtrates were blown dry with a stream of nitrogen. The residue was purified by high performance liquid chromatography over Purospher Star RP-18 (20 g, 5 μ m; eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃OH/CH₃CN (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10-20 min) 75/25/0). The desired fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with CH₂Cl₂/Na₂CO₃ solution. The extract was purified through ExtrelutTM and the organic phase was blown dry with a stream of nitrogen. The residue was dried under vacuum at 60°C, yielding 0.008 g of 2',4'-difluoro-N-[4-[4-[2-oxo-1-phenyl-2-(propylamino)ethyl] -1-piperidiny]phenyl]-[1,1'-biphenyl]-2-carboxamide identified as compound No. 84 in the following table F-1. Compound identified as No. 68 in the following table F-1 was similarly prepared while using the same experimental procedure.

20 EXAMPLE B.18

The resin of example A.19 (0.0001 mole) was washed three times with dioxane. 1,4-Dioxane (3 ml) was added. 2-Methylphenylboronic acid (0.0008 mole) in dioxane (1 ml) was added. KOH (2 M) (0.8 ml) was added. The mixture was shaken for 30 minutes under argon atmosphere. PdCl₂(PPh₃)₂ (0.00002 mole) in NMP (0.5 ml) was added. The mixture was shaken for two hours at 96°C. Again, PdCl₂(PPh₃)₂ in 0.5 ml of NMP was added and the mixture was warmed for two hours at 96°C. The mixture was allowed to cool to room temperature, then filtered and the filter residue was washed three times with DMF, three times with H₂O, three times with DMF, three times with methanol, three times with DCM, three times with methanol, and three times with methanol. A mixture of TFA/DCM/TIS (4 ml) was added. The mixture was shaken for 60 minutes at room temperature, filtered, washed with a mixture of TFA/DCM/TIS (2 ml) and DCM (2 ml), and the filtrates were blown dry with a stream of nitrogen. The residue was purified by high performance liquid chromatography over Purospher Star RP-18 (20 g, 5 μ m; eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃OH/CH₃CN (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10-20 min) 75/25/0), yielding 0.004 g of N-[4-[4-[2-(ethylamino)-2-

oxo-1-phenylethyl]-1-piperidinyl]phenyl]-2'-methyl-[1,1'-biphenyl]-2-carboxamide identified as compound No. 85 in the following table F-1.

Compounds identified as No. 86 to No. 96 in the following table F-1 were similarly prepared while using the same experimental procedure.

EXAMPLE B.19

Argon gas was bubbled through a mixture of the resin of example A.23 (0.0001 mole) and 3,5-dichlorobenzeneboronic acid (0.0008 mole) in K_2CO_3 (2M in H_2O) (0.0008 mole) and 1,4-dioxane (5 ml) for five minutes. Palladium (II)

acetate (0.00001 mole) in dioxane (0.5 ml) was added and the reaction mixture was warmed and shaken for sixteen hours at 97°C, then cooled, filtered, washed with DMF (three times), water (three times), DMF (three times), then three times with firstly methanol and secondly DCM. A mixture of TFA/DCM/TIS (4 ml, 5/93/2) was added and the mixture was shaken for one hour then filtered.

A mixture of TFA/DCM/TIS (2 ml, 5/93/2) was added. The mixture was shaken for 10 minutes, then filtered, washed with DCM (3 ml) and the filtrate was blown dry under nitrogen at 50°C. The residue was purified by HPLC over Purospher Star RP-18-e (20 g, 5 μm ; eluent: ((0.5% NH_4OAc in H_2O)/ CH_3CN 90/10)/ CH_3OH/CH_3CN (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min)

0/0/100, (18.10-20 min) 75/25/0). The desired fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with CH_2Cl_2 /aqueous K_2CO_3 solution. The extract was purified through ExtrelutTM and the organic phase was blown dry with a stream of nitrogen. The residue was dried under vacuum at 60 °C, yielding 0.008 g of 3',5'-dichloro-N-[4-[4-[2-oxo-1-phenyl-2-(propylamino)ethyl]-1-piperidinyl]phenyl] -[1,1'-biphenyl]-2-carboxamide identified as compound No. 97 in the following table F-1.

Compounds identified as No. 42 to 51, 53, 82, 83, and 98 to 118 in the following table F-1 were similarly prepared while using the same experimental procedure.

EXAMPLE B.20

a) A suspension of 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.086 g, 0.00014 mole) in NMP (1 ml) was added to the resin of example A.38 (0.2 g, 0.00014 mole) and sodium tertbutoxide (0.242 g, 0.00252 mole). Homopiperazine (0.126 g, 0.0021 mole) in NMP (2 ml) was added and the mixture was stirred under argon. Tris(dibenzylidene acetone) di-palladium (0.026 g, 0.000028 mole) in NMP (1 ml) was added and the reaction mixture was shaken for 19 hours at 105°C. The mixture was cooled, filtered and the filter residue was washed with DMF, water, DMF (three times), H_2O (three

times), DMF (three times), CH₃OH (three times), CH₂Cl₂ (three times), CH₃OH (three times) and NMP (two times). NMP (3 ml) was added.

b) Methyl-bromophenylacetate (0.16 g, 0.0007 mole) in NMP (1 ml) was added to the product obtained in step (a). DIPEA (0.3 ml) was added and the mixture was shaken for 18 hours at room temperature. The mixture was filtered, washed with DMF and water, then with DMF (three times), water (three times), DMF (three times), methanol (three times), DCM (three times), methanol (three times) DCM (three times).

TFA/TIS/CH₂Cl₂ (49/2/49) (4 ml) was added and the mixture was shaken for one hour at room temperature. The mixture was filtered and more TFA/TIS/CH₂Cl₂ (49/2/49) (1.5 ml) was added. The mixture was shaken for 15 minutes, filtered, washed with DCM (2 ml), then the filtrates were blown dry under nitrogen. The residue was purified by high performance liquid chromatography over Purospher Star RP-18 (20 g, 5 μ m; eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃CN/CH₃OH (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10 – 20 min) 75/25/0). The desired fractions were collected and the organic solvent was evaporated. The aqueous concentrate was treated with an aqueous sodium carbonate solution, then extracted with DCM. The extract was separated through ExtrelutTM and the filtrates were blown dry under nitrogen at 50°C, yielding 0.021 g of the compound identified as compound 119 in the following table F-1.

Compounds identified as No. 120 to No. 128 in the following table F-1 were similarly prepared while using the same experimental procedure.

Example B.21

A mixture of intermediate (17) (0.019 mol) and Na₂CO₃ (0.019 mol) in DMF (125ml) was stirred at room temperature. Methyl α -bromo- α -phenylacetate (0.01907 mol) was added dropwise. The mixture was stirred for 3 hours. The solvent was evaporated. The residue was taken up in water and DCM. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 100/0; 99.5/0.5). The pure fractions were collected and the solvent was evaporated yielding a residue that separated in its enantiomers by high performance liquid chromatography over Chiralpak AD (eluent : hexane/ethanol 70/30). The desired fractions were collected and the solvent was evaporated, yielding, after crystallisation from 2-propanol, compound (229), mp. 158°C, $[\alpha]_D^{20} = -28.86^\circ$ (c = 24.95 mg/5 ml in CH₃OH); and

compound (230), mp. 160°C, $[\alpha]_D^{20} = +27.69^\circ$ (c = 24.95 mg/5 ml in CH₃OH).

Example B.22

Methyl α -bromo- α -phenylacetate (0.0010 mole) in NMP (1 ml) was added to
5 resin (IX-a) (0.0002 mol) in NMP (3 ml). *N,N*-diisopropylethylamine (0.0023
mol) was added and the reaction mixture was shaken for 48 hours at room
temperature, then filtered and the filter residue was washed with DMF (3 times),
water (3 times), DMF (3 times), methanol (3 times), DCM (3 times), methanol
(3 times) and DCM (3 times). A mixture of TFA/TIS/CH₂Cl₂ (49/2/49) (4 ml) was
10 added. the reaction mixture was shaken for 2 hours at room temperature, then
filtered and again a mixture of TFA/TIS/CH₂Cl₂ (49/2/49) (4 ml) was added.
The reaction mixture was shaken for another 15 minutes, then filtered and the
filtrates were blown dry under nitrogen. The residue was purified by high
performance liquid chromatography over Purospher Star RP-18 (20 g, 5 μ m;
15 eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃CN/CH₃OH (0 min) 75/25/0,
(10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10 – 20 min) 75/25/0). The
desired fractions were collected and the organic solvent was evaporated. The
aqueous concentrate was treated with an aqueous sodium carbonate solution,
then extracted with DCM. The extract was separated through Extrelut™ and
20 the filtrates were blown dry under nitrogen at 50°C, yielding compound (184) in
the following table F-1.

Example B.23

Dimethylallyl alcohol (0.00017 mol) was added to a mixture of α -phenyl-4-[4-
25 [[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino] phenyl]-1-
piperidineacetic acid monohydrochloride (as prepared in Example B.16.a)
(0.000084 mol) in DCM (4 ml) and *N,N*-diisopropylethylamine (0.00010 mol)
stirred under nitrogen at -20°C and the mixture was stirred for 10 minutes.
1-[Bis(dimethylamino)methylene]-tetrafluoroborate(1-)-1*H*-benzotriazolium
30 3-oxide (TBTU) (0.00013 mol) was added and the reaction mixture was stirred
for 30 minutes at -20°C. The mixture was allowed to gradually warm to room
temperature and the reaction mixture was stirred for 75 hours at room
temperature. The reaction mixture was washed with water (1 ml), then filtered
through Extrelut™ and the filter residue was rinsed three times with 3 ml of
35 DCM. The filtrate was evaporated and the residue was purified by HPLC
(Waters column, with Xterra MS C18; eluent: [(0.5% NH₄OAc in H₂O)/CH₃CN
90/10)]/CH₃OH/CH₃CN (0 min) 75/25/0, (10 min) 0/50/50, (16 min) 0/0/100,
(18.10-20.00 min) 75/25/0). The product fractions were collected and the
organic solvent was evaporated. The aqueous concentrates were partitioned

between DCM and an aqueous Na_2CO_3 solution. The combined organic layers were separated, dried, filtered and the filtrate was blown dry under nitrogen at 50°C . The residue was dried (vacuum, 60°C), yielding compound (220).

5 Example B.24

N,N-diisopropylethylamine (0.0010 mol) was added to a mixture of α -phenyl-4-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-1-piperidine-acetic acid monohydrochloride (as prepared in Example B.16.a) (0.000084 mol) in DCM (4 ml). Ethanol (0.00017 mol) and 1-[bis(dimethylamino)methylene]-
10 tetrafluoroborate(1-)-1*H*-benzotriazolium 3-oxide (TBTU) (0.00013 mol) were added. The reaction mixture was stirred for 75 hours at room temperature. Water (1 ml) was added. The mixture was stirred for 30 minutes, then filtered through ExtrelutTM, rinsed 3 times with DCM (3 ml each time) and the filtrate was evaporated. The residue was dissolved in DCM, washed with 1 ml of 1 N
15 HCl, filtered through ExtrelutTM and the filtrate was washed with a saturated aqueous NaHCO_3 solution (1 ml). This mixture was filtered through ExtrelutTM. The filtrate was collected, and the filter residue was washed with DCM (2 x 4 ml). The filtrate was evaporated. Each residue was purified by HPLC (Waters column, with Xterra MS C18; eluent: [(0.5% NH_4OAc in H_2O)/ CH_3CN
20 90/10)]/ $\text{CH}_3\text{OH}/\text{CH}_3\text{CN}$ (0 min) 75/25/0, (10 min) 0/50/50, (16 min) 0/0/100, (18.10-20.00 min) 75/25/0). The product fractions were collected and the organic solvent was evaporated. The aqueous concentrates were partitioned between DCM and an aqueous Na_2CO_3 solution. The combined organic layers were separated, dried, filtered and the filtrate was blown dry under nitrogen at
25 50°C . The residue was dried (vacuum, 60°C), yielding compound (222).

Example B.25

- a) A mixture of compound (39) (0.0014 mol) in concentrated HCl (25 ml) and dioxane (20 ml) was stirred and refluxed for 4 hours, cooled and poured out into
30 water. The mixture was extracted with DCM. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was triturated in DIPE. The precipitate was filtered off and dried, yielding 0.48 g of α -phenyl-1-[4-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]-4-piperidineacetic acid (intermediate 26) (mp. 196°C).
- 35 b) Ethylbromide (1.2 equivalent, 0.00010 mol) was added to a mixture of intermediate (26) (0.000084 mol) in DMF (5 ml) and Cs_2CO_3 (0.00018 mol) and the reaction mixture was stirred for 3 hours at 70°C . The solvent was evaporated. The residue was partitioned between water and DCM. The extract's solvent was evaporated. The residue was purified by HPLC (Waters

column, with Xterra MS C18; eluent: [(0.5% NH₄OAc in H₂O)/CH₃CN 90/10)]/CH₃CN (0 min) 85/15, (10 min) 10/90, (16 min) 0/100, (18.10-20.00 min) 85/15). The product fractions were collected and the organic solvent was evaporated. The aqueous concentrates were extracted and the extract's solvent was evaporated, yielding compound (243).

Example B.26

Acetyl chloride (0.0007 mol) was added to resin (XI) (0.00011 mol) in DCM (4 ml). *N,N*-dimethyl-4-pyridinamine (0.00011 mol) was added. *N,N*-diisopropyl-ethylamine (0.0011 mol) was added and the reaction mixture was shaken overnight at room temperature. The mixture was filtered and the filter residue was washed with DCM, methanol, DCM, methanol, CH₂Cl₂, methanol, and DCM. TFA/TIS/CH₂Cl₂ (49/2/49) (4 ml) was added and the mixture was shaken for 2 hours at room temperature. The mixture was filtered, More TFA/TIS/CH₂Cl₂ (49/2/49) (2 ml) was added and the mixture was shaken for 15 minutes, then filtered and the filter residue was washed with DCM (2 ml). The filtrates were blown dry under nitrogen. The residue was purified by HPLC over Purospher Star RP-18 (20 g, 5 μm; eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃OH/CH₃CN (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10-20 min) 75/25/0). The desired fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted and the extract's solvent was evaporated, yielding 0.001 g of compound (253).

Example B.27

Methanol (0.5 ml) was added to resin (XII) (0.0002 mol) in DCM (4 ml). *N,N*-dimethyl-4-pyridinamine (0.0002 mol) was added. DIPEA (0.002 mol) was added and the reaction mixture was shaken for 18 hours at room temperature. The mixture was filtered and the filter residue was washed with DCM (3 x), methanol (3 x), DCM (3 x), methanol (3 x), DCM (3 x), methanol (3 x), DCM (3 x). TFA/TIS/CH₂Cl₂ (49/2/49) (4 ml) was added and the mixture was shaken for 2 hours at room temperature. The mixture was filtered, More TFA/TIS/CH₂Cl₂ (49/2/49) (2 ml) was added and the mixture was shaken for 15 minutes, then filtered and the filter residue was washed with DCM (2 ml). The filtrates were blown dry under nitrogen. The residue was purified by HPLC over Purospher Star RP-18 (20 g, 5 μm; eluent: ((0.5% NH₄OAc in H₂O)/CH₃CN 90/10)/CH₃OH/CH₃CN (0 min) 75/25/0, (10.00 min) 0/50/50, (16.00 min) 0/0/100, (18.10-20 min) 75/25/0). The desired fractions were collected and the

organic solvent was evaporated. The aqueous concentrate was extracted and the extract's solvent was evaporated, yielding 0.002 g of compound (251).

Example B.28

- 5 A mixture of intermediate (31) (0.0006 mol), ethylamine hydrochloride (0.0015 mol), EDCI (0.0007 mol), HOBT (0.0007 mol) and triethylamine (0.0015 mol) in DCM (10 ml) was stirred at room temperature overnight. Water was added. The mixture was extracted with DCM. The organic layer was separated, dried, filtered, and the solvent was evaporated. The residue was
10 purified by column chromatography over kromasil (eluent : DCM). The pure fractions were collected and the solvent was evaporated, yielding 0.034 g of compound (276).

Example B.29











- 15 A mixture of intermediate (29) (0.0045 mol), 3-thienyl boronic acid (0.036 mol), $\text{PdCl}_2(\text{PPh}_3)_2$ (0.0009 mol) and Na_2CO_3 (0.072 mol) in dioxane (50 ml) was stirred and refluxed for 30 minutes. Water was added. The mixture was extracted with ethyl acetate. The organic layer was separated, dried, filtered, and the solvent was evaporated. The residue was purified by column
20 chromatography over silica gel (eluent : DCM/ethyl acetate 90/10). The pure fractions were collected and the solvent was evaporated, yielding compound (267) (mp. 150°C).













Example B.30













- A mixture of intermediate (29) (0.0025 mol), 3-pyridinyl boronic acid (0.02 mol),
25 $\text{PdCl}_2(\text{PPh}_3)_2$ (0.005 mol) and Na_2CO_3 (0.04 mol) in dioxane (30ml) was stirred and refluxed for 3 hours. Water was added. The mixture was extracted with ethyl acetate. The organic layer was separated, dried, filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 97/3/0.1), yielding compound (270)
30 (mp. 194°C).













- Table F-1 lists the polyarylcarboxamide compounds of the present invention, together with their detailed formulae, that were prepared according to one of the above examples B.1 to B.20. In this table, the abbreviation ". $\text{C}_2\text{HF}_3\text{O}_2$ " stands for the trifluoroacetate salt, ". $\text{C}_3\text{H}_8\text{O}$ " stands for the 2-propanolate salt,
35 and ". CH_4O " stands for the methanolate salt of the said compound.













Table F-1













	
Co. No. 1; Ex. B.1; mp. 208°C	Co. No. 38; Ex. B.7; .C ₂ HF ₃ O ₂
	
Co. No. 40; Ex. B.9; mp. 143°C	Co. No. 81; Ex. B.14; mp. 195°C
	
Co. No. 68; Ex. B.17	Co. No. 54; Ex. B.12; mp. 123°C
	
Co. No. 55; Ex. B.12	Co. No. 56; Ex. B.12
	
Co. No. 57; Ex. B.12	Co. No. 58; Ex. B.12













	
Co. No. 59 ; Ex. B.12	Co. No. 60; Ex. B.12
	
Co. No. 61 ; Ex. B.12; mp. 142°C	Co. No. 62 ; Ex. B.13; mp. 151°C
	
Co. No. 3; Ex. B.2	Co. No. 2; Ex. B.2
	
Co. No. 4; Ex. B.2	Co. No. 5; Ex. B.2
	
Co. No. 6 ; Ex. B.2	Co. No. 7; Ex. B.2
	
Co. No. 8; Ex. B.2	Co. No. 9; Ex. B.2













	
Co. No. 10; Ex. B.2	Co. No. 11; Ex. B.2; mp. 218°C
	
Co. No. 13 ; Ex. B.3	Co. No. 12; Ex. B.3
	
Co. No. 14; Ex. B.3	Co. No. 15; Ex. B.3
	
Co. No. 16; Ex. B.3	Co. No. 17; Ex. B.3
	
Co. No. 18; Ex. B.3	Co. No. 19; Ex. B.3
	
Co. No. 20; Ex. B.3	Co. No. 22 ; Ex. B.4













	
Co. No. 21; Ex. B.4	Co. No. 23; Ex. B.4
	
Co. No. 24; Ex. B.4	Co. No. 25; Ex. B.4
	
Co. No. 26; Ex. B.4	Co. No. 27; Ex. B.4
	
Co. No. 28; Ex. B.4	Co. No. 30; Ex. B.5
	
Co. No. 29; Ex. B.5	Co. No. 31; Ex. B.5
	
Co. No. 32; Ex. B.5	Co. No. 33; Ex. B.5













	
Co. No. 34 ; Ex. B.5	Co. No. 35; Ex. B.5
	
Co. No. 36; Ex. B.5	Co. No. 63; Ex. B.15
	
Co. No. 64; Ex. B.15	Co. No. 65; Ex. B.15
	
Co. No. 66; Ex. B.15	Co. No. 67 ; Ex. B.15
	
Co. No. 37; Ex. B.6; .C ₂ HF ₃ O ₂	Co. No. 41; Ex. B.10; mp. 138°C
	
Co. No. 84; Ex. B.17	Co. No. 97; Ex. B.19















	
Co. No. 82; Ex. B.19	Co. No. 83; Ex. B.19
	
Co. No. 42; Ex. B.19	Co. No. 43; Ex. B.19
	
Co. No. 44; Ex. B.19	Co. No. 45; Ex. B.19
	
Co. No. 46; Ex. B.19	Co. No. 47; Ex. B.19
	
Co. No. 48; Ex. B.19	Co. No. 49; Ex. B.19
	
Co. No. 52; Ex. B.11	Co. No. 85; Ex. B.18













	
Co. No. 86; Ex. B.18	Co. No. 87; Ex. B.18
	
Co. No. 88; Ex. B.18	Co. No. 89; Ex. B.18
	
Co. No. 90; Ex. B.18	Co. No. 91; Ex. B.18
	
Co. No. 92; Ex. B.18	Co. No. 93; Ex. B.18
	
Co. No. 94; Ex. B.18	Co. No. 95; Ex. B.18
	
Co. No. 96; Ex. B.18	Co. No. 69; Ex. B.16













	
Co. No. 70; Ex. B.16	Co. No. 71; Ex. B.16
	
Co. No. 72; Ex. B.16	Co. No. 73; Ex. B.16
	
Co. No. 74; Ex. B.16	Co. No. 75; Ex. B.16
	
Co. No. 76; Ex. B.16	Co. No. 77; Ex. B.16
	
Co. No. 78; Ex. B.16	Co. No. 79; Ex. B.16
	
Co. No. 80; Ex. B.16	Co. No. 39; Ex. B.8











	
Co. No.50 ; Ex. B.19	Co. No. 51; Ex. B.19
	
Co. No. 53; Ex. B.19	Co. No. 98; Ex. B.19
	
Co. No. 99; Ex. B.19	Co. No. 100; Ex. B.19
	
Co. No. 101; Ex. B.19	Co. No. 102; Ex. B.19
	
Co. No. 103; Ex. B.19	Co. No. 104; Ex. B.19
	
Co. No. 105; Ex. B.19	Co. No. 106; Ex. B.19













	
Co. No. 107; Ex. B.19	Co. No. 108; Ex. B.19
	
Co. No. 109; Ex. B.19	Co. No. 110; Ex. B.19
	
Co. No. 111; Ex. B.19	Co. No. 112; Ex. B.19
	
Co. No. 113; Ex. B.19	Co. No. 114; Ex. B.19
	
Co. No. 115; Ex. B.19	Co. No. 116; Ex. B.19
	
Co. No. 117; Ex. B.19	Co. No. 118; Ex. B.19













	
Co. No. 119; Ex. B.20	Co. No. 120; Ex. B.20
	
Co. No. 121; Ex. B.20	Co. No. 122; Ex. B.20
	
Co. No. 123; Ex. B.20	Co. No. 124; Ex. B.20
	
Co. No. 125; Ex. B.20	Co. No. 126; Ex. B.20
	
Co. No. 127; Ex. B.20	Co. No. 128; Ex. B.20
	
Co. No. 129; Ex. B.3	Co. No. 130; Ex. B.19
	
Co. No. 131; Ex. B.17	Co. No. 132; Ex. B.17













	
Co. No. 133; Ex. B.17	Co. No. 134; Ex. B.18
	
Co. No. 135; Ex. B.18	Co. No. 136; Ex. B.17
	
Co. No. 137; Ex. B.17	Co. No. 138; Ex. B.17
	
.CH ₄ O (1:1); Co. No. 139; Ex. B.9; mp. 152°C	Co. No.140; Ex. B.9; mp. 231°C
	
Co. No. 141; Ex. B.9	Co. No. 142; Ex. B.9
	
Co. No. 143; Ex. B.9	Co. No. 144; Ex. B.9; mp. 138°C













	
Co. No. 145; Ex. B.9; mp. 217°C	Co. No. 146; Ex. B.9; mp. 136°C
	
Co. No. 147; Ex. B.9 ; mp. 132°C	.C ₃ H ₈ O (1:1); Co. No. 148; Ex. B.9
	
Co. No. 149; Ex. B.9; mp. 125°C	.HCl (1:2); Co. No. 150; Ex. B.9; mp. 151°C
	
Co. No. 151; Ex. B.9	Co. No. 152; Ex. B.9
	
Co. No. 153; Ex. B.9	Co. No. 154; Ex. B.9
	
Co. No. 155; Ex. B.9	Co. No. 156; Ex. B.9













	
Co. No. 157; Ex. B.8; mp. 174°C	HCl (1:2) .H ₂ O (1:1); Co. No. 158; Ex. B.9
	
Co. No. 159; Ex. B.22	Co. No. 160; Ex. B.22
	
Co. No. 161; Ex. B.9	Co. No. 162; Ex. B.9
	
Co. No. 163; Ex. B.9	Co. No. 164; Ex. B.9
	
Co. No. 165; Ex. B.9	Co. No. 166; Ex. B.9

	
Co. No. 167; Ex. B.9	Co. No. 168; Ex. B.9
	
Co. No. 169; Ex. B.9	Co. No. 170; Ex. B.9
	
Co. No. 171; Ex. B.9	Co. No. 172; Ex. B.12
	
Co. No. 173; Ex. B.12	Co. No. 174; Ex. B.12
	
Co. No. 175; Ex. B.12	Co. No. 176; Ex. B.12
	
Co. No. 177; Ex. B.12	Co. No. 178; Ex. B.12











	
Co. No. 179; Ex. B.12	Co. No. 180; Ex. B.16; mp. 192°C
	
Co. No. 181; Ex. B.16; mp. 142°C	.HCl (1:1); Co. No. 182; Ex. B.9
	
.HCl (1:1); Co. No. 183; Ex. B.9	Co. No. 184; Ex. B.22
	
Co. No. 185; Ex. B.22	Co. No. 186; Ex. B.22
	
Co. No. 187; Ex. B.22	Co. No. 188; Ex. B.22
	
Co. No. 189; Ex. B.22	Co. No. 190; Ex. B.22













	
Co. No. 191; Ex. B.22	Co. No. 192; Ex. B.22
	
Co. No. 193; Ex. B.22	Co. No. 194; Ex. B.22
	
Co. No. 195; Ex. B.22	Co. No. 196; Ex. B.22
	
Co. No. 197; Ex. B.22	Co. No. 198; Ex. B.22
	
Co. No. 199; Ex. B.22	Co. No. 200; Ex. B.22
	
Co. No. 201; Ex. B.22	Co. No. 202; Ex. B.22











	
.HCl (1:1) .C ₃ H ₈ O (1:1); Co. No. 203; Ex. B.8	Co. No. 204; Ex. B.22
	
Co. No. 205; Ex. B.22	Co. No. 206; Ex. B.22
	
Co. No. 207; Ex. B.22	Co. No. 208; Ex. B.22
	
Co. No. 209; Ex. B.22	Co. No. 210; Ex. B.22
	
Co. No. 211; Ex. B.22	Co. No. 212; Ex. B.22
	
Co. No. 213; Ex. B.22	Co. No. 214; Ex. B.22









	
Co. No. 215; Ex. B.22	Co. No. 216; Ex. B.22
	
Co. No. 217; Ex. B.22	Co. No. 218; Ex. B.22
	
Co. No. 219; Ex. B.22	Co. No. 220; Ex. B.23
	
Co. No. 221; Ex. B.23	Co. No. 222; Ex. B.24
	
Co. No. 223; Ex. B.24	Co. No. 224; Ex. B.24
	
Co. No. 225; Ex. B.24	Co. No. 226; Ex. B.24

Co. No. 227; Ex. B.24	Co. No. 228; Ex. B.24
(-); Co. No. 229; Ex. B.21; mp. 158°C; [α] _D ²⁰ = -28.86° (c = 24.95 mg/5 ml in CH ₃ OH)	(+); Co. No. 230; Ex. B.21; mp. 160°C; [α] _D ²⁰ = +27.69° (c = 24.38 mg/5 ml in CH ₃ OH)
Co. No. 231; Ex. B.17	Co. No. 232; Ex. B.17
Co. No. 233; Ex. B.17	Co. No. 234; Ex. B.17
Co. No. 235; Ex. B.17	Co. No. 236; Ex. B.17

	
Co. No. 237; Ex. B.17	Co. No. 238; Ex. B.17
	
(A); Co. No. 239; Ex. B.9	(B); Co. No. 240; Ex. B.9
	
(A); Co. No. 241; Ex. B.8	(B); Co. No. 242; Ex. B.8
	
Co. No. 243; Ex. B.25	Co. No. 244; Ex. B.25
	
Co. No. 245; Ex. B.25	Co. No. 246; Ex. B.25

	
Co. No. 247; Ex. B.25	Co. No. 248; Ex. B.25
	
Co. No. 249; Ex. B.25	Co. No. 250; Ex. B.25
	
Co. No. 251; Ex. B.27	Co. No. 252; Ex. B.27
	
Co. No. 253; Ex. B.26	Co. No. 254; Ex. B.2
	
Co. No. 255; Ex. B.2	Co. No. 256; Ex. B.2
	
Co. No. 257; Ex. B.2	Co. No. 258; Ex. B.9

	
Co. No. 259; Ex. B.9	Co. No. 260; Ex. B.9
	
Co. No. 261; Ex. B.9	Co. No. 262; Ex. B.9
	
Co. No. 263; Ex. B.9	(-); Co. No. 264; Ex. B.24; mp. 147°C; [α] _D ²⁰ = -18.03° (c = 25.51 mg/5 ml in CH ₃ OH)
	
(+); Co. No. 265; Ex. B.24; mp. 142°C; [α] _D ²⁰ = +15.76° (c = 25.70 mg/5 ml in CH ₃ OH)	Co. No. 266; Ex. B.24
	
Co. No. 267; Ex. B.29; mp. 150°C	Co. No. 268; Ex. B.9; mp. 114°C

	
Co. No. 269; Ex. B.29; mp. 90°C	Co. No. 270; Ex. B.30; mp. 194°C
	
Co. No. 271; Ex. B.29; mp. 60°C	Co. No. 272; Ex. B.28; mp. 114°C
	
Co. No. 273; Ex. B.30	Co. No. 274; Ex. B.28
	
Co. No. 275; Ex. B.28	Co. No. 276; Ex. B.28

C. Pharmacological examples.

C.1. Quantification of the secretion of ApoB.

HepG2 cells were cultured in 24-well plates in Minimal Essential Medium Rega
 3 containing 10 % fetal calf serum. Rega 3 has the following composition: CaCl₂
 5 (264 µg/ml), KCl (400 µg/ml), MgSO₄·7H₂O (200 µg/ml), NaCl (6800 µg/ml),
 NaHCO₃ (850 µg/ml), NaH₂PO₄·H₂O (158 µg/ml), D-glucose (1000 µg/ml),
 phenol red (10 µg/ml), L-alanine (8.9 µg/ml), L-arginine HCl (12 µg/ml),
 L-asparagine·H₂O (15 µg/ml), L-aspartic acid (13.3 µg/ml), L-cystine (24 µg/ml),
 10 L-glutamic acid (14.7 µg/ml), glycine (7.5 µg/ml), L-histidine·HCl·H₂O
 (42 µg/ml), L-isoleucine (52 µg/ml), L-leucine (52 µg/ml), L-lysine·HCl (72.5
 µg/ml), L-methionine (15 µg/ml), L-phenylalanine (32 µg/ml), L-proline (11.5
 µg/ml), L-serine (10.5 µg/ml), L-threonine (48 µg/ml), L-tryptophan (10 µg/ml),
 L-tyrosine (36 µg/ml), L-valine (46 µg/ml), D-Ca pantothenate (1 µg/ml), choline
 15 chloride (1 µg/ml), folic acid (1 µg/ml), 1-enositol (2 µg/ml), nicotinamide

(1 µg/ml), pyridoxal HCl (1 µg/ml), riboflavin (0.1 µg/ml) and thiamine HCl (1 µg/ml).

At 70% confluency, the medium was changed and the test compound or carrier (dimethylsulfoxide, 0.4 % final concentration) was added. After 24 hours of incubation, the medium was transferred to Eppendorf tubes and cleared by centrifugation. A sheep antibody directed against either apo B was added to the supernatant and the mixture was kept at 8°C for 24 hours. Then, rabbit anti-sheep antibody was added and the immune complex was allowed to precipitate for 24 hours at 8°C. The immunoprecipitate was pelleted by centrifugation for 25 minutes at 1320 g and washed twice with a buffer containing 40 mM 4-morpholinepropanesulfonic acid, 40 mM NaH₂PO₄, 100 mM NaF, 0.2 mM dithiothreitol, 5 mM ethylenediamine tetraacetic acid, 5 mM ethylenebis (oxyethylenenitrilo) tetraacetic acid, 1 % Triton-X-100, 0.5 % sodium deoxycholate, 0.1 % sodium dodecylsulphate, 0.2 µM leupeptin and 0.2 µM phenylmethylsulphonylfluoride. Radioactivity in the pellet was quantified by liquid scintillation counting.

Resulting IC₅₀ values are enumerated in Table C.1 for a number of compounds No. 1 to No. 123.

TABLE C.1

Co. No.	pIC ₅₀
1	8.103
2	6.974
3	7.591
4	5.523
5	6.802
6	6.967
7	6.583
8	7.221
9	6.655
10	6.618
11	8.335
12	6.636
13	7.523
14	5.523
15	6.688
16	7.077
17	6.702

Co. No.	pIC ₅₀
18	6.687
19	6.578
20	6.005
21	6.611
22	6.984
23	5.523
24	6.072
25	6.542
26	6.561
27	5.885
28	6.115
29	5.523
30	6.782
31	5.894
32	5.621
33	5.523
34	5.809

Co. No.	pIC ₅₀
35	5.523
36	7.523
37	7.507
38	8.179
39	7.791
40	7.586
41	7.666
42	5.523
43	5.751
44	5.981
45	5.523
46	6.336
47	6.702
48	6.198
49	6.627
50	7.028
51	7.163

Co. No.	pIC ₅₀
52	6.531
53	8.736
54	8.103
55	7.523
56	7.523
57	8.121
58	7.523
59	7.523
60	7.523
61	8.414
62	7.19
63	6.912
64	6.799
65	6.62
66	7.099
67	6.608
68	7.523
69	8.051
70	7.523
71	7.987
72	7.523
73	8.216
74	7.523
75	7.943
76	7.286
77	7.523
78	7.488
79	7.301
80	6.448
81	6.749
82	7.011
83	7.364
84	7.1
85	6.888
86	7.075
87	8.688

Co. No.	pIC ₅₀
88	7.523
89	6.44
90	7.851
91	8.061
92	7.199
93	7.141
94	7.356
95	7.523
96	7.493
97	6.63
98	7.237
99	7.523
99	8.062
100	7.935
101	6.684
102	7.732
103	7.133
104	8.1
105	7.043
106	6.6
107	6.535
108	6.725
109	5.833
110	6.8
111	6.655
112	6.363
113	6.938
114	6.078
115	5.766
116	6.414
117	6.916
118	6.895
119	6.757
120	6.447
121	5.676
122	6.383

Co. No.	pIC ₅₀
123	6.618
129	5.557
130	6.444
131	6.38
132	7.299
133	< 5.523
134	7.185
135	6.826
136	6.829
137	5.752
138	7.003
139	7.065
140	7.693
141	7.601
142	6.944
143	6.631
144	6.695
145	6.732
146	6.467
147	6.542
148	7.219
149	7.38
150	6.761
151	6.213
152	7.025
153	5.809
154	5.634
156	6.915
157	6.97
158	7.671
159	6.973
160	7.489
161	7.162
162	7.015
163	6.602
164	7

Co. No.	pIC ₅₀
165	7.482
166	7.444
167	> 7.523
168	6.881
174	7.035
175	> 7.523
176	6.873
177	6.583
178	7.465
179	6.395
180	6.945
181	8.48
182	8.118
183	7.666
184	8.505
185	7.312
186	6.698
187	7.386
188	8
189	6.979
190	8.193
191	8.143
192	6.802
193	6.629
194	7.367
195	7.047
196	6.999
197	7.783
198	7.136
200	7.096
201	> 7.523

Co. No.	pIC ₅₀
202	6.736
203	8.333
204	6.369
205	7.323
206	6.106
207	> 7.523
208	8.126
209	8.099
210	6.316
211	7.702
212	> 7.523
213	> 7.523
214	> 7.523
215	> 7.523
216	7.327
217	5.57
218	> 7.523
219	7.105
220	> 7.523
221	> 7.523
222	7.716
223	6.432
224	6.254
225	7.17
227	6.727
229	7.409
230	8.31
231	> 7.523
232	> 7.523
233	> 7.523
234	> 7.523

Co. No.	pIC ₅₀
235	> 7.523
236	7.663
237	> 7.523
238	7.824
239	7.855
240	7.982
241	7.578
242	8.133
243	> 7.523
245	6.28
246	6.762
247	< 5.523
248	7.164
249	< 5.523
250	6.632
251	8.016
252	7.915
253	6.887
254	7.541
255	8.298
256	7.821
257	7.857
258	6.856
259	6.377
260	> 7.523
261	6.129
262	6.253
263	6.555
266	7.591

C.2. MTP assay

MTP activity was measured using an assay similar to one described by J.R. Wetterau and D.B. Zilversmit in *Chemistry and Physics of Lipids* (1985) 38, 205-222. In order to prepare the donor and acceptor vesicles, the appropriate lipids in chloroform were put into a glass test tube and dried under a stream of

nitrogen. A buffer containing 15 mM Tris-HCl (pH 7.5), 1 mM ethylenediamine tetra-acetic acid, 40 mM NaCl, 0.02 % NaN₃ (assay buffer) was added to the dried lipid. The mixture was vortexed briefly and the lipids were then allowed to hydrate for 20 minutes on ice. Vesicles were then prepared by bath sonication (using a Branson 2200 device) at room temperature for at most 15 minutes. Butylated hydroxytoluene was included in all vesicle preparations at a concentration of 0.1%. The lipid transfer assay mixture contained donor vesicles (40 nmole phosphatidylcholine, 7.5 mole % cardiolipin and 0.25 mole % glycerol tri [1-¹⁴C]-oleate), acceptor vesicles (240 nmol phosphatidylcholine) and 5 mg bovine serum albumin in a total volume of 675 μ l in a 1.5 ml microcentrifuge tube. Test compounds were added dissolved in dimethylsulfoxide (0.13 % final concentration). After 5 minutes of pre-incubation at 37°C, the reaction was started by the addition of MTP in 100 μ l of a dialysis buffer. The reaction was stopped by the addition of 400 μ l diethylaminoalkyl (DEAE)-52 cellulose (Sephadex) pre-equilibrated in 15 mM Tris-HCl (pH 7.5), 1 mM ethylenediamine tetra-acetic acid and 0.02% NaN₃ (1:1 volume/volume). The mixture was agitated for 4 minutes and centrifuged for 2 minutes at maximum speed in an Eppendorf centrifuge (4°C) to pellet the DEAE-52-bound donor vesicles. An aliquot of the supernatant containing the acceptor liposomes was counted and the [¹⁴C]-counts were used to calculate the percent triglyceride transfer from donor to acceptor vesicles. Resulting IC₅₀ values are enumerated in Table C.2 for some of the above-referred compounds.

TABLE C.2

Co. No.	pIC ₅₀
1	7.864
3	7.735
6	6.703
7	6.44
10	< 5.523
11	8.136
16	6.682
39	7.922
40	8.344
41	8.063


Co. No.	pIC ₅₀
54	8.269
55	8.37
57	8.163
58	7.799
60	8.082
61	8.32
62	7.98
75	8.077
87	8.495
199	8.334

Co. No.	pIC ₅₀
203	8.682
222	8.075
229	7.279
230	8.439
232	7.602
239	8.703
241	7.985
242	7.94
264	7.497
265	8.25

CLAIMS

1. Polyarylcaboxamide compounds of formula (I)

(48)




the N-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

- Z_1 is selected from $(CH_2)_n$ wherein n is 1 to 3, CH_2CH_2O and OCH_2CH_2 ;
- 10 - Z_2 is $(CH_2)_m$ wherein m is 1 or 2;
- X_1 represents O, CH_2 , CO, NH, CH_2O , OCH_2 , CH_2S , SCH_2 or a direct bond;
- X_2 and X_3 are each independently selected from CH, N and a sp^2 carbon atom;
- R_1 is hydrogen or C_{1-4} alkyl;
- 15 - Ar^1 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one or two R_3 substituents;
- Ar^2 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl,
- 20 - pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one, two or three R_4 substituents;
- each R_2 and R_3 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo and trifluoromethyl;
- 25 - each R_4 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo, hydroxy, mercapto, cyano, nitro, C_{1-4} alkylthio or polyhalo C_{1-6} alkyl, amino, C_{1-4} alkylamino and di(C_{1-4} alkyl)amino;
- p^1 and p^2 are each 0 to 2;
- p^3 is 0 to 3;
- 30 - X_1 and R_4 taken together with the aromatic rings Ar^1 and Ar^2 to which they are attached may form a fluoren-1-yl or a fluoren-4-yl group;

- A represents a C₁₋₆alkanediyl substituted with one or two groups selected from aryl, heteroaryl and C₃₋₁₀cycloalkyl; or when X₃ is CH, A may also represent a nitrogen atom substituted with hydrogen, C₁₋₁₀alkyl, aryl, heteroaryl, arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl or C₃₋₁₀cycloalkyl;
 - 5 - B represents hydrogen; C₁₋₁₀alkyl; aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkoxycarbonyl, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₆alkyl; heteroarylC₁₋₁₀alkyl;
 - 10 C₃₋₁₀cycloalkyl; polyhaloC₁₋₆alkyl; C₃₋₆alkenyl; C₃₋₆alkynyl; NR₆R₇; or OR₈;
 - R₆ and R₇ each independently represent hydrogen, C₁₋₁₀alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl;
 - 15 arylC₁₋₁₀alkyl, heteroarylC₁₋₁₀alkyl, C₃₋₁₀cycloalkyl, C₇₋₁₀polycycloalkyl, polyhaloC₁₋₆alkyl, C₃₋₈alkenyl, C₃₋₈alkynyl, fused benzo-C₅₋₈cycloalkyl, and wherein R₆ and R₇ taken together with the nitrogen atom to which they are attached may form a C₄₋₈ saturated heterocyclic radical;
 - R₈ represents C₁₋₁₀alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C₁₋₄alkyloxy, amino, C₁₋₁₀alkylamino, di(C₁₋₁₀alkyl)amino, C₁₋₁₀acyl, C₁₋₁₀alkylthio, C₁₋₁₀alkylaminocarbonyl and di(C₁₋₁₀alkyl)aminocarbonyl; arylC₁₋₁₀alkyl; heteroarylC₁₋₁₀alkyl; and
 - 20 C₃₋₁₀cycloalkyl; C₇₋₁₀polycycloalkyl; polyhaloC₁₋₆alkyl; C₃₋₈alkenyl; C₃₋₈alkynyl; or fused benzo-C₅₋₈ cycloalkyl.
- 25
2. Polyarylcarboxamide compounds according to claim 1, wherein Z₁, Z₂, X₂ and X₃ taken together form a six-membered heterocycle.
 3. Polyarylcarboxamide compounds according to claim 1 or claim 2, wherein
 - 30 Z₁, Z₂, X₂ and X₃ taken together form a piperidine or piperazine group and X₁ is a direct bond.
 4. Polyarylcarboxamide compounds according to any of claims 1 to 3, wherein R₂ and R₃ are each hydrogen and R₄ is hydrogen, trifluoromethyl, chloro or


tert-butyl.

5. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4, wherein an intermediate phenylene amine having the formula (II)



(II)


- wherein Z_1 , Z_2 , X_2 , X_3 , p^1 , R_1 , R_2 , A and B are as defined in formula (I), is reacted with a polyarylcboxylic acid or halide having the formula (III),




(III)

- wherein X_1 , Ar^1 , Ar^2 , p^2 , p^3 , R_3 and R_4 are as defined in formula (I) and Y_1 is selected from hydroxy and halo, in at least one reaction-inert solvent and optionally in the presence of a suitable base, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.

6. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4 wherein X_3 is nitrogen, wherein an intermediate having the formula (IV)




wherein Z_1 , Z_2 , X_1 , X_2 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I) and X_3 is nitrogen, is reacted with a reactant having the formula (V)




wherein A and B are as defined in formula (I) and Y_2 is selected from halo, tosyloxy, mesyloxy, naphthylsulfonyloxy, or $-A-Y_2$ is R'_5COR_5 , wherein R_5 and R'_5 are such that the radical R'_5CHR_5 is encompassed by the definition of A in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one suitable nucleophilic substitution activator and/or a suitable base, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.

7. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4, wherein an intermediate having the formula




wherein Z_1 , Z_2 , X_1 , X_2 , X_3 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 , R_4 and A are as defined in formula (I) and Y_3 is selected from halo and hydroxy, is reacted with a reactant of the formula BH , wherein B is NR_6R_7 or OR_8 and R_6 , R_7 and R_8 are as defined in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one suitable coupling reagent and/or a suitable base, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.

8. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4 wherein X_3 is nitrogen and wherein A is a group suitable for a Michael addition reaction, wherein an intermediate having the formula




(IV)

- wherein Z_1 , Z_2 , X_1 , X_2 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I) and X_3 is nitrogen, is reacted with a reactant of the formula (VII)




(VII)

wherein B is as defined in formula (I), and Y_4 and Y'_4 are such that the




is encompassed by the definition of A in formula (I), in at least one reaction-inert solvent, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.

9. A process for preparing a polyarylcarboxamide compound according to any of claims 1 to 4, wherein an intermediate having the formula (VIII)



(VIII)


- 5 wherein X_1 , p^1 , p^2 , p^3 , Ar^1 , Ar^2 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I) and Y_5 is selected from halo, $B(OH)_2$, alkylboronates and cyclic analogues thereof, is reacted with a reactant having the formula (IX)



(IX)


- 10 wherein Z_1 , Z_2 , X_3 , A and B are as defined in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one transition metal coupling reagent and/or at least one suitable ligand, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.
- 15 thereof.

10. A process for preparing a polyarylcarboxamide compound according to any of claims 1 to 4, wherein an intermediate having the formula (X)



(X)


- 20 wherein p^1 , p^2 , p^3 , Ar^1 , Ar^2 , X_1 , R_1 , R_2 , R_3 and R_4 are as defined in formula (I), is reacted with a reactant having the formula (XI)



(XI)


- wherein Z_1 , Z_2 , X_3 , A and B are as defined in formula (I), one of Y_6 and Y_7 is selected from bromo, iodo and trifluoromethylsulfonate and the other of Y_6 and Y_7 is selected from tri(C_{1-4} alkyl) tin, $B(OH)_2$, alkylboronates and cyclic analogues thereof, in at least one reaction-inert solvent and optionally in the presence of at least one transition metal coupling reagent and/or at least one suitable ligand, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.

11. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4, wherein an intermediate having the formula (XII)




(XII),

- wherein Z_1 , Z_2 , X_2 , X_3 , p^1 , p^2 , Ar^1 , R_1 , R_2 , R_3 , A and B are as defined in formula (I) and Y_8 is selected from bromo, iodo and trifluoromethylsulfonate, is reacted either with an arylboric acid having the formula (XIII a)



(XIII-a)

- wherein p^3 , Ar^2 and R_4 are as defined in formula (I), or with an aryl-tin reactant having the formula (XIII b)




(XIII b)


wherein R_3 , R_4 , Ar^2 and p^3 are as defined in formula (I), in at least one reaction-inert solvent and optionally in the presence of at least one transition metal coupling reagent and/or at least one suitable ligand,, the said process further optionally comprising converting a compound of formula (I) into an addition salt thereof, and/or preparing or stereochemically isomeric forms thereof.

12. A process for preparing a polyarylcaboxamide compound according to any of claims 1 to 4, wherein B is NR_6R_7 , from a polyarylcaboxamide compound according to claim 1 wherein B is OR_5 , comprising in a first step hydrolyzing the latter and in a second step reacting the resulting corresponding carboxylic acid with an amine having the formula HNR_6R_7 in at least one reaction-inert solvent and further optionally comprising converting the resulting compound of formula (I) wherein B is NR_6R_7 into an addition salt thereof, and/or preparing stereochemically isomeric forms thereof.


13. Compounds of any of the formulae :




(II)



(IV)



(VI)



(XII)

5

wherein:

- Z_1 is selected from $(CH_2)_n$ wherein n is 1 to 3, CH_2CH_2O and OCH_2CH_2 ;
- Z_2 is $(CH_2)_m$ wherein m is 1 or 2;
- 10 - X_1 represents O, CH_2 , CO, NH, CH_2O , OCH_2 , CH_2S , SCH_2 or a direct bond;
- X_2 and X_3 are each independently selected from CH, N and a sp^2 carbon atom;
- R_1 is hydrogen or C_{1-4} alkyl;
- Ar^1 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one or two R_3 substituents;
- 15 - Ar^2 is an aromatic ring selected from phenyl, naphthalenyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolyl, furanyl and thienyl, optionally substituted with one, two or three R_4 substituents;
- 20 - each R_2 and R_3 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo and trifluoromethyl;

- each R_4 is independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, halo, hydroxy, mercapto, cyano, nitro, C_{1-4} alkylthio or polyhalo C_{1-6} alkyl, amino, C_{1-4} alkylamino and di(C_{1-4} alkyl)amino;
- p^1 and p^2 are each 0 to 2;
- 5 - p^3 is 0 to 3;
- X_1 and R_4 taken together with the aromatic rings Ar^1 and Ar^2 to which they are attached may form a fluoren-1-yl or a fluoren-4-yl group;
- A represents a C_{1-6} alkanediyl optionally substituted with one or two groups selected from aryl, heteroaryl and C_{3-10} cycloalkyl; oxygen; or a direct bond;
- 10 - B represents hydrogen; C_{1-10} alkyl; aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, nitro, C_{1-4} alkyloxy, amino, C_{1-10} alkylamino, di(C_{1-10} alkyl)amino, C_{1-10} acyl, C_{1-10} alkylthio, C_{1-10} alkoxycarbonyl, C_{1-10} alkylaminocarbonyl and di(C_{1-10} alkyl)aminocarbonyl; aryl C_{1-10} alkyl; heteroaryl C_{1-10} alkyl; C_{3-10} cycloalkyl;
- 15 polyhalo C_{1-6} alkyl; C_{3-6} alkenyl; C_{3-6} alkynyl; NR_6R_7 ; or OR_8 ;
- R_6 and R_7 each independently represent hydrogen, C_{1-10} alkyl, aryl or heteroaryl each optionally substituted with a group selected from halo, cyano, C_{1-4} alkyloxy, amino, C_{1-10} alkylamino, di(C_{1-10} alkyl)amino, C_{1-10} acyl, C_{1-10} alkylthio, C_{1-10} alkylaminocarbonyl and di(C_{1-10} alkyl)aminocarbonyl;
- 20 aryl C_{1-10} alkyl, heteroaryl C_{1-10} alkyl, C_{3-10} cycloalkyl, C_{7-10} polycycloalkyl, polyhalo C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, fused benzo- C_{5-8} cycloalkyl, and wherein R_6 and R_7 taken together with the nitrogen atom to which they are attached may form a C_{4-8} saturated heterocyclic radical;
- R_8 represents C_{1-10} alkyl, aryl or heteroaryl each optionally substituted with a
- 25 group selected from halo, cyano, nitro, C_{1-4} alkyloxy, amino, C_{1-10} alkylamino, di(C_{1-10} alkyl)amino, C_{1-10} acyl, C_{1-10} alkylthio, C_{1-10} alkylaminocarbonyl and di(C_{1-10} alkyl)aminocarbonyl; aryl C_{1-10} alkyl; heteroaryl C_{1-10} alkyl; C_{3-10} cycloalkyl; C_{7-10} polycycloalkyl; polyhalo C_{1-6} alkyl; C_{3-8} alkenyl; C_{3-8} alkynyl; or fused benzo- C_{5-8} cycloalkyl;
- 30 - when X_3 is CH, A may also represent a nitrogen atom substituted with hydrogen, C_{1-10} alkyl, aryl, heteroaryl, aryl C_{1-10} alkyl, heteroaryl C_{1-10} alkyl or C_{3-10} cycloalkyl;
- Y_1 and Y_3 are each independently selected from hydroxy and halo;

- Y₅ is selected from halo, B(OH)₂, alkylboronates and cyclic analogues thereof; and
 - Y₆ and Y₈ are each independently selected from bromo, iodo and trifluoromethylsulfonate,
- 5 useful as intermediates for the preparation of polyarylcaboxamide compounds according to any of claims 1 to 4.
- 10 14. A pharmaceutical composition comprising at least one pharmaceutically acceptable carrier and a therapeutically effective amount of a compound according to any of claims 1 to 4.
- 15 15. A polyarylcaboxamide compound according to any of claims 1 to 4 for use as a medicine.
- 15 16. A polyarylcaboxamide compound according to claim 13 for use in the treatment of hyperlipidemia, obesity, type II diabetes, atherosclerosis, ischemic heart disease, peripheral vascular disease, cerebral vascular disease, hypercholesterolemia, hypertriglyceridemia, pancreatitis or coronary artery disease.
- 20 17. A pharmaceutical composition according to claim 12, further comprising at least one additional lipid-lowering agent.
- 25 18. A method of treating a condition selected from hyperlipidemia, obesity, type II diabetes, atherosclerosis, ischemic heart disease, peripheral vascular disease, cerebral vascular disease, hypercholesterolemia, hypertriglyceridemia, pancreatitis or coronary artery disease, comprising administering to a mammal in need of such treatment a polyarylcaboxamide compound according to any of claims 1 to 4 in an
- 30 amount sufficient to decrease the secretion of apolipoprotein B.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number
WO 02/20501 A3

(51) International Patent Classification⁷: C07D 211/34,
295/14, 213/40, A61K 31/495, A61P 3/06

(21) International Application Number: PCT/EP01/09926

(22) International Filing Date: 27 August 2001 (27.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00203067.4 4 September 2000 (04.09.2000) EP

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Patent
Department, Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MEERPOEL,
Lieven [BE/BE]; Janssen Pharmaceutica N.V., Turn-
houtseweg 30, B-2340 Beerse (BE). ROEVENS, Peter,
Walter, Maria [BE/BE]; Janssen Pharmaceutica N.V.,
Turnhoutseweg 30, B-2340 Beerse (BE). BACKX, Leo,
Jacobus, Jozef [BE/BE]; Janssen Pharmaceutica N.V.,
Turnhoutseweg 30, B-2340 Beerse (BE). VAN DER
VEKEN, Louis, Jozef, Elisabeth [BE/BE]; Janssen
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse
(BE). VIELLEVOYE, Marcel [NL/BE]; Janssen Phar-
maceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Agent: VERBERCKMOES, Filip; Patent Department,
Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340
Beerse (BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE,
SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL,
SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE,
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR),
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only


Published:

- with international search report

(88) Date of publication of the international search report:
27 June 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: POLYARYLCARBOXAMIDES USEFUL AS LIPID LOWERING AGENTS



(57) Abstract: Polyarylcarboxamide compounds of formula (I) are useful as lipid lowering agents.

WO 02/20501 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/09926

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D211/34 C07D295/14 C07D213/40 A61K31/495 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 32582 A (DAUGAN ALAIN CLAUDE MARIE ; GLAXO GROUP LTD (GB)) 8 June 2000 (2000-06-08) intermediates page 23, line 4 -page 26, line 30 claims 1,24-26	1-18
A	WO 96 40640 A (QUALLICH GEORGE J ; DORFF PETER H (US); CHANG GEORGE (US); PFIZER ()) 19 December 1996 (1996-12-19) cited in the application abstract example 163	1,14-16, 18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

22 February 2002

Date of mailing of the international search report

06/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Diederer, J

INTERNATIONAL SEARCH REPORT

In tional Application No

PCT/EP 01/09926

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 27979 A (SQUIBB BRISTOL MYERS CO) 2 July 1998 (1998-07-02) cited in the application abstract examples 11-74	1,14-16, 18
A	US 5 760 246 A (TINO JOSEPH A ET AL) 2 June 1998 (1998-06-02) cited in the application abstract example 429	1,14-16, 18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/09926

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0032582	A	08-06-2000	AU 1656600 A	19-06-2000
			BR 9915895 A	21-08-2001
			CN 1334808 T	06-02-2002
			CZ 20011973 A3	12-09-2001
			WO 0032582 A1	08-06-2000
			EP 1135378 A1	26-09-2001
			NO 20012688 A	31-05-2001
WO 9640640	A	19-12-1996	CA 2223574 A1	19-12-1996
			HU 9601566 A2	29-09-1997
			WO 9640640 A1	19-12-1996
			AU 3585399 A	16-09-1999
			AU 703493 B2	25-03-1999
			AU 5478496 A	19-12-1996
			BG 62442 B1	30-11-1999
			BG 100637 A	31-03-1997
			BR 9602628 A	08-09-1998
			CN 1141918 A ,B	05-02-1997
			CZ 9601644 A3	15-01-1997
			EP 0832069 A1	01-04-1998
			FI 974440 A	27-01-1998
			HR 960270 A1	31-12-1997
			KR 225713 B1	15-10-1999
			LV 11615 A	20-12-1996
			LV 11615 B	20-04-1997
			NO 962385 A	09-12-1996
			NZ 286733 A	26-02-1998
			PL 314636 A1	09-12-1996
			RO 116897 B1	30-07-2001
			RU 2141478 C1	20-11-1999
			SG 44952 A1	19-12-1997
			SI 9600183 A	30-04-1997
			SK 72696 A3	05-11-1997
			TR 961058 A2	21-12-1996
			US 5919795 A	06-07-1999
WO 9827979	A	02-07-1998	AU 5513298 A	17-07-1998
			WO 9827979 A1	02-07-1998
			US 6281228 B1	28-08-2001
			US 5965577 A	12-10-1999
US 5760246	A	02-06-1998	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.